

Phases of Gauge Theories

Michael C. Ogilvie*

Dept. of Physics, Washington University, St. Louis, MO 63130 USA

Abstract

One of the most fundamental questions we can ask about a given gauge theory is its phase diagram. In the standard model, we observe three fundamentally different types of behavior: QCD is in a confined phase at zero temperature, while the electroweak sector of the standard model combines Coulomb and Higgs phases. Our current understanding of the phase structure of gauge theories owes much to the modern theory of phase transitions and critical phenomena, but has developed into a subject of extensive study. After reviewing some fundamental concepts of phase transitions and finite-temperature gauge theories, we discuss some recent work that broadly extends our knowledge of the mechanisms that determine the phase structure of gauge theories. A new class of models with a rich phase structure has been discovered, generalizing our understanding of the confinement-deconfinement transition in finite-temperature gauge theories. Models in this class have space-time topologies with one or more compact directions. On $R^3 \times S^1$, the addition of double-trace deformations or periodic adjoint fermions to a gauge theory can yield a confined phase in the region where the S^1 circumference L is small, so that the coupling constant is small, and semiclassical methods are applicable. In this region, Euclidean monopole solutions, which are constituents of finite-temperature instantons, play a crucial role in the calculation of a non-perturbative string tension. We review the techniques used to analyze this new class of models and the results obtained so far, as well as their application to finite-temperature phase structure, conformal phases of gauge theories and the large- N limit.

*mco@physics.wustl.edu

I. INTRODUCTION: GAUGE THEORIES

Gauge theories are central to the modern understanding of the fundamental forces of nature, and form the foundation of the standard model of particle physics. Gauge theories promote global symmetries, such as the $U(1)$ invariance associated with conservation of electric charge, to local symmetries. In the case of quantum electrodynamics (QED), the quantum field theory of electromagnetic interactions, local symmetry requires the introduction of a new vector field, associated with the photon. The standard model of particle physics is based on a local symmetry group of the form $SU(2) \times U(1) \times SU(3)$. The standard model contains the Weinberg-Salam-Glashow model of the electroweak interactions, with gauge group $SU(2) \times U(1)$, which unifies the electromagnetic interaction with the weak interaction. The number of gauge particles equals the rank of the gauge group, in this case four: the massless photon γ and the massive vector boson $W^\pm$ and Z^0 . The other part of the standard model is quantum chromodynamics (QCD), with gauge group $SU(3)$, describes the strong interactions of quarks. There are eight gauge fields in QCD, describing the gluons that bind quarks inside of hadrons. It is widely believed that the standard model is not a complete description of particle physics, and there are many different proposals for what lies beyond the standard model. Gauge theories are integral to all such proposals, even for those where four-dimensional gauge theories are low-energy effective theories, as in string theory.

One of the most fundamental questions we can ask about a given gauge theory is its phase structure. In the standard model, we observe three fundamentally different types of behavior: the familiar Coulomb behavior associated with the massless photon; the Higgs mechanism, responsible for the masses of the $W^\pm$ and Z^0 ; and the confinement of quarks and gluons by the gluon fields. These properties are characteristics of different phases: QCD is in a confined phase at zero temperature, while the electroweak sector of the standard model combines Coulomb and Higgs phases. The phase structure of gauge theories has been extensively studied. A large part of the interest in this subject centers on QCD at non-zero temperature and density, but there is now great interest in the phase structure of gauge theories associated with physics beyond the standard model.

It is often convenient theoretically to view QCD in a simplified way, in a form without dynamical quarks; this is often referred to as pure $SU(3)$ gauge theory. As a consequence

of dimensional transmutation, the dimensionless gauge coupling g^2 can be replaced by a parameter Λ with dimensions of energy. Key observables with units of mass such as hadron masses become pure numbers times Λ . In other words, the pure gauge theory is a theory with no adjustable dimensionless parameters, making it difficult to carry out analytical approximations. Including dynamical quarks with realistic masses does not make analytical work easier. However, there are many parameters that can be varied within the context of gauge theories, shedding light on important aspects of gauge theory behavior. Aside from their great experimental interest, finite-temperature gauge theories offer a dimensionless parameter T/Λ which can be used, for example, to change QCD from a confining phase at low temperatures to a plasma phase at high temperatures. This additional parameter tells us something about the physics of both phases. It is also natural to study the phase structure of gauge as other parameters, such as the number of colors N or the number of quark flavors N_f , are varied.

Recent work has shown the existence of a new class of gauge theory models with important and desirable properties. All of these models have one or more compact directions, and the most developed case is the geometry $R^3 \times S^1$, which is the geometry of Euclidean gauge theories at finite temperature when the circumference L of S^1 is identified with the inverse temperature $\beta = 1/T$. In addition to L , there are many parameters that can be used to examine the phase structure. Unlike conventional finite-temperature gauge theories, this new class can be put into a confined phase when $L \ll \Lambda$ and $g^2(L) \ll 1$ [1]. Euclidean monopoles, the constituents of finite-temperature instantons, are essential to a semiclassical calculation of the string tension in this region. Moreover, this small- L phase is smoothly connected to the conventional, large- L confining phase [2]. There is a price to be paid for this. In the case of models with adjoint fermions, the use of periodic boundary conditions in the compact direction removes the spectral positivity of the transfer matrix in the compact direction, so L cannot be identified with β and must be regarded as a spatial direction. Models with double-trace deformations can be motivated as the heavy-quark limit of periodic adjoint fermions, and have a similar problem. The gains, however, are great: an analytic understanding of confinement in a class of four-dimensional models, the discovery of many new phases, and new approaches to conformality and to the large- N limit.

The analysis of this new class of models uses techniques and ideas from many areas of theoretical physics, and the results are broadly interesting as well. In section II, we

review the modern theory of phase transitions and critical phenomena, concentrating on key concepts and their appearance in a field-theoretic context. Section III introduces the basics of finite-temperature gauge theories, including important symmetries of gauge theories associated with confinement and chiral symmetry breaking. Sections IV through VI discuss recent developments. Section IV discussed the phase structure of gauge theories on $R^3 \times S^1$, an arena where it has proven possible to demonstrate confinement in certain models using semiclassical methods. Section V focuses on the general issue of gauge theory phase structure in the context of conformality and duality in gauge theories. In section VI, $SU(N)$ gauge theories in the large- N limit are considered, and promising new approaches are compared with older formulations. A final section concludes and summarizes.

The notations used throughout are as follows: All field theories are taken to be in Euclidean space unless noted otherwise. Lower-case Greek indices are used for space-time and the metric in Euclidean space is

$$g_{\mu\nu} = g^{\mu\nu} = \delta_{\mu\nu}. \quad (1)$$

Roman indices in the range $j \cdots n$ generally denote “spatial” directions on $R^3 \times S^1$, i.e., the three directions orthogonal to the compact direction. Roman indices in the range $a \cdots d$ generally label group generators, while capital letters are used to denote group representations: F, Adj, S, A, R . S^k is the k -dimensional surface of a $(k+1)$ -dimensional hypersphere, so S^1 is the unit circle. T^k is the k -dimensional hypertorus, so T^1 is also S^1 .

II. MODERN THEORY OF PHASE TRANSITIONS AND CRITICAL PHENOMENA

A. Order parameters and symmetries

The modern theory of phase transitions begins with Landau-Ginsburg theory [3], and the key concepts of order parameters and free energies as functions of those order parameters. The simplest examples of order parameters come from ferromagnetic spin systems, where the magnetization $m_j(\vec{x})$ is the order parameter, with j specifying the number of components. In order for the magnetization to be treated as a continuous variable, it is necessary to imagine that some smoothing procedure be applied to average the microscopic spins over many unit cells. The magnetization is naturally a three-component vector, but underly-

ing microscopic physics may cause it to be effectively one- or two-dimensional. Assuming macroscopic isotropy, the magnetization has a natural internal symmetry group $O(3)$, $O(2)$ or $Z(2)$ depending on whether the order parameter is three, two or one dimensional. Thus the dimensionality of the order parameter need not be the same as the dimensionality of the system. Suppose for a moment we are interested in the free energy of a spin system in three spatial dimensions with a uniform magnetization ϕ which we take to be a scalar. This is the case where the magnetization has an easy axis; then ϕ is the projection of the magnetization onto the easy axis. The key assumption of Landau-Ginsburg theory is that the free energy may be written as an integral over a local free energy density $f(\phi)$ which is an analytic function of ϕ as well as any underlying parameters, including the temperature. Thus the free energy is given by

$$F[\phi] = \int d^3x f(\phi) = \int d^3x \left[\frac{r}{2}\phi^2 + \frac{\lambda}{4}\phi^4 + \dots \right] \quad (2)$$

where only terms even in ϕ are allowed by the $Z(2)$ symmetry. If higher-order terms in the polynomial expansion are to be ignored, then we must have $\lambda > 0$ for stability. If $r > 0$, the free energy is minimized when $\phi = 0$. As shown in figure 1, if $r < 0$, the free energy will be minimized when

$$\phi = \pm \sqrt{\frac{-r}{\lambda}} \quad (3)$$

giving the most common example of the idea of spontaneously broken symmetry. The free energy concept can be generalized to that of a function of a spatially-dependent magnetization $\phi(\vec{x})$. Including gradient terms in the expansion of the free energy, we obtain

$$F[\phi] = \int d^3x \left[\frac{a}{2} (\nabla\phi)^2 + \frac{r}{2}\phi^2 + \frac{\lambda}{4}\phi^4 + \dots \right] \quad (4)$$

where higher-order terms in $\nabla\phi$ are suppressed. It is now possible to imagine calculating the free energy in an ensemble with an external field, via the functional integral

$$Z[h] = \int [d\phi] \exp \left[-\frac{1}{T} F[\phi] + \int d^3x h(\vec{x}) \phi(\vec{x}) \right] \quad (5)$$

This is a three-dimensional Euclidean scalar field theory, the simplest example of the confluence of quantum field theory (at zero temperature) and classical statistical mechanics. Within Landau-Ginsburg theory, the point $r = 0$ marks the location of a second-order phase transition, where the correlation length $\xi \propto r^{-1/2}$ becomes infinite. From a field-theoretic point of view, the inverse of the correlation length is a mass. From the modern point of view, a second-order phase transition is intrinsically associated with a massless excitation.




Figure 1: The Landau-Ginsburg free energy density $f(\phi)$ for the case $r < 0$ where the symmetry $\phi \rightarrow -\phi$ is spontaneously broken.

B. Universality

Landau-Ginsburg theory explains how non-analyticities arise in the vicinity of second-order critical point. For example, the critical point can be determined as a function of temperature: $r(T_C) = 0$ locates the critical point. Analyticity of r in T gives $r \propto (T - T_c)$ near the critical point and hence $\phi(T) \propto \sqrt{T_c - T}$ in the low-temperature, symmetry-broken phase. The non-analytic behavior is captured in a set of critical indices, which characterize the critical behavior. For example, $\phi(T) \propto (T_c - T)^\beta$ so the critical index β is $1/2$ in Landau-Ginsburg theory. Landau-Ginsburg theory leads to a simple universality: models with the same Landau-Ginsburg theory have the same phase diagram near the critical point, when parametrized in terms of the Landau-Ginsburg parameters, and the critical exponents at second-order phase transitions are identical. This universality was supported by the known exact equivalence between the Ising model and a lattice binary alloy model, and led to the identification of the liquid-gas critical end point with the similar critical point in spin systems. However, both theoretical and experimental determinations of critical indices led to the conclusion that that critical indices, while falling into universal classes, were not given by the Landau-Ginsburg values [4]. Furthermore, it was clear that the spatial dimensionality d of the system must play a role. For example, classical spin systems with $d = 1$ with short-range interactions do not have phase transitions except at $T = 0$, in contradiction to the predictions of Landau-Ginsburg theory. It was clear that fluctuations, ignored in Landau-Ginsburg theory, were crucial. This led to the development of the modern renormalization

group [5].

C. Renormalization group

The renormalization group takes many forms, but common to all approaches is the idea of a fixed point in the space of couplings. Consider an Ising model of spatial dimension d . It has two parameters: the lattice spacing a and a dimensionless coupling J which is the energy of a ferromagnetic bond divided by $k_B T$. The Hamiltonian is given by a sum over nearest-neighbor interactions:

$$-\frac{\mathcal{H}}{k_B T} = \sum_{nn} J \sigma_j \sigma_k \quad (6)$$

where the spins σ_j are associated with lattice sites j and take on the values ± 1 . On general grounds, the correlation length ξ between spins must have the form $\xi = aF(J)$. At the critical point J_c , the correlation length is infinite: $F(J_c) = \infty$. If an exact or approximate relation between ξ and J is known, we can define a simple renormalization group transformation as a change in the lattice spacing by a scale factor $s > 1$ such that $a \rightarrow a' = sa$ and $J \rightarrow J'(J)$ with the correlation length ξ fixed: $\xi' = \xi$. In other words, we must have $sF(J') = F(J)$ or

$$J' = F^{-1}(s^{-1}F(J)) \quad (7)$$

At the critical point J_c , we must have $J' = J$, which is a fixed point of this renormalization group transformation. There are two equivalent points of view: From a physical point of view the correlation length is constant, but from a lattice point of view the dimensionless correlation length $F(J)$, measured in units of lattice spacing, decreases away from J_c . In the Ising model with $d \geq 2$, there are actually three fixed points. There are two fixed points with $\xi = 0$, at $J = 0$ and at $J = \infty$, corresponding to $T = 0$ and $T = \infty$ respectively. The second-order phase transition is associated with the non-trivial fixed point between them.

It is generally necessary to introduce many parameters into the effective Hamiltonian describing the system at a length scale λa , which can be denoted as the set $\{J_a\}$. In Ising models, such additional couplings include a next-nearest-neighbor coupling and a four-spins around a square coupling. It is here that the renormalization group has its explanatory power. In the vicinity of a fixed point $J^* = \{J_a^*\}$, the renormalization group transform

which changes the scale by a factor λ can be linearized as

$$J'_a - J_a^* \simeq \sum_b R_{ab} (J_b - J_b^*). \quad (8)$$

The matrix R can be diagonalized and the corresponding linear combinations of spin interactions identified; the latter are usually referred to as operators and denoted $\mathcal{O}_c$. In the new basis for interactions the Hamiltonian can be written as $\mathcal{H} = \sum_c J_c \mathcal{O}_c$ and a renormalization group transformation has the form $J'_c - J_c^* = s^{p_c} (J_c - J_c^*)$. If $p_c > 0$, then the renormalization group flow generated by $J \rightarrow J'$ will take J_c away from its fixed point value. In this case, the coupling and its corresponding operators are termed relevant. On the other hand, if $p_c < 0$, the renormalization group flow will naturally take J_c towards its fixed point value, at least within some basin of attraction where the linearization of the renormalization group is valid. Such couplings and operators are called irrelevant. There is also the case $p_c = 0$, the marginal case, where higher-order effects in the renormalization group transformation must be considered. In the simplest case, a second-order phase transition is associated with the existence of one relevant coupling J_r , accompanied by a potentially infinite set of irrelevant couplings. This situation occurs, for example, in the Ising model in the absence of symmetry-breaking fields. Within the linear approximation, the hyper-plane in coupling space defined by $J_r = J_r^*$, is the critical surface where the correlation length is infinite. More generally, the set of renormalization group trajectories that flow into the fixed point forms a critical surface of codimension one in parameter space. The existence of irrelevant variables explains universality: many different models lie in the same universality class, and their critical behavior is determined not by the particular microscopic model, but by the physics near the fixed point.

In the field-theoretic approach to critical phenomena, regularization of ultraviolet divergences typically requires the introduction of a parameter μ with dimensions of mass. Typical regularization schemes include a simple ultraviolet cutoff, Pauli-Villars regularization and dimensional regularization. The observed coupling constants are defined in terms of the arbitrary scale μ , and are therefore called running coupling constants. A change in μ does not change the underlying physics, but implies a corresponding change in the coupling constants, because a change in conventions does not change the physics of the system. In the case of a pure gauge theory, there is only one coupling $g(\mu)$, and its running is described by

$$\mu \frac{dg}{d\mu} = \beta(g) \quad (9)$$

where $\beta(g)$ is the renormalization group beta function. In pure $SU(N)$ gauge theories, the perturbative β function is negative, indicating that the renormalization group flow is towards $g = 0$ at large μ . We say that $g = 0$ is an ultraviolet fixed point, or that the theory is asymptotically free. There is an important connection here with the construction of the continuum limit of lattice gauge theories, where the inverse of lattice spacing a plays the role of an ultraviolet regulator. Physical quantities, such as a glueball mass M , cannot depend on the lattice spacing, but can be written parametrically as

$$M = a^{-1} F(g(a)) \tag{10}$$

where $g(a)$ is the coupling constant defined in the lattice action. The continuum limit, where M is to remain finite as $a \rightarrow 0$, can only be taken by simultaneously adjusting g such that $F(g) \rightarrow 0$. From the perspective of fixed lattice spacing, the continuum limit can only be taken at a second-order critical point, and physical masses become small in lattice units. Although the most common gauge theories have a simple fixed point structure in four dimensions, non-trivial fixed points play an important role in some scenarios for physics beyond the standard model, as will be discussed in section V.

The close connection between quantum field theories at zero temperature in d Euclidean dimensions with classical statistical mechanics models was crucial in developing the field-theoretic approach to critical phenomena. The simplest example of this connection is between the simple model of ferromagnetism described above and the ϕ^4 field theory. There is a simple rule of thumb which is helpful in matching up field-theoretic properties with renormalization group properties: a super-renormalizable interaction is relevant, a renormalizable interaction is marginal, and a non-renormalizable interaction is irrelevant. For example, in the case of the $\lambda\phi^4$ field theory in $d = 4 - \epsilon$ dimensions [6–8], the ϕ^4 interaction is marginal in four dimensions, with an infrared fixed point at $\lambda = 0$. For $\epsilon > 0$, a non-trivial infrared fixed point $\lambda^* > 0$ is present, and the ϕ^2 term is the only relevant operator near the critical point. From this point of view, Landau-Ginsburg theory fails to give the correct critical indices for $d < 4$ because it is essentially an expansion around $\lambda = 0$ rather than λ^* . Above four dimensions, the IR fixed point is at $\lambda = 0$, and the critical indices are correctly predicted by Landau-Ginsburg theory. The field theoretic approach allows a precise statement of universality: starting from the most general renormalizable field theory using the order parameter as the field, the second-order phase transitions and their properties follow from

the fixed points of the theory. Thus, the universality class of a second-order phase transition depends on the dimensionality of space and the symmetry group of the order parameter.

Although the renormalization group has led to tremendous progress in our understanding of second-order phase transitions, the vast majority of phase transitions observed in nature are first-order. Such transitions are generally accompanied by a discontinuity in one or more order parameter, and there is a latent heat associated with the transition. Within the framework of the effective potential, first-order transitions occur when there are two or more disconnected global minima of the free energy. A simple model that illustrates the general concept has a Landau-Ginsburg free energy density of the form

$$f(\phi) = \frac{r}{2}\phi^2 + \frac{\lambda}{4}\phi^4 + \frac{g}{6}\phi^6 \quad (11)$$

Unlike the simpler ϕ^4 theory, with $g > 0$ to ensure stability we need not have $\lambda > 0$. For $\lambda > 0$, there is a second-order phase transition at $r = 0$. However, if $r < 0$, there is a first-order phase transition at $\lambda = -4\sqrt{rg/3}$ where two non-trivial minima have the same free energy as the local minimum at $\phi = 0$.

This model contains an important lesson: Although the behavior of systems at second-order transitions is determined by the local structure of the free energy density $f(\phi)$, the overall phase structure is determined by the location of the global minimum of the free energy. This is usually more difficult to obtain in systems with several order parameters and complicated interactions.

D. The effective potential

The close correspondence between classical statistical mechanics and quantum field theory extends beyond critical phenomena. In thermodynamics and statistical mechanics, different independent variables, free energies and statistical ensembles are often used. These ideas, so useful in statistical physics, have direct analogs in quantum field theory. We begin with the case of quantum field theories at zero temperature, and later extend our results to finite temperature. Consider the following generalization of the partition function

$$Z[J] = \int [d\phi] \exp \left[-S[\phi] + \int d^{d+1}x J(x) \phi(x) \right] \quad (12)$$

This is a generating functional for all n -point correlation functions.

$$\frac{Z[J]}{Z[0]} = 1 + \sum_{n=1}^{\infty} \frac{1}{n!} \int d^{d+1}x_1 \dots d^{d+1}x_n J(x_1) \dots J(x_n) \langle \phi(x_1) \dots \phi(x_n) \rangle. \quad (13)$$

The logarithm of $Z[J]$, $-W[J] = \log Z[J]$ is the generating functional of connected correlation functions

$$-W[J] = -W[0] + \sum_{n=1}^{\infty} \frac{1}{n!} \int d^{d+1}x_1 \dots d^{d+1}x_n J(x_1) \dots J(x_n) \langle \phi(x_1) \dots \phi(x_n) \rangle_c \quad (14)$$

where the connected correlation functions are, for example,

$$\langle \phi(x_1) \phi(x_2) \rangle_c = \langle \phi(x_1) \phi(x_2) \rangle - \langle \phi(x_1) \rangle \langle \phi(x_2) \rangle \quad (15)$$

While $W[0]$ is typically not physically meaningful, the difference $W[J] - W[0]$ can be interpreted in the case of constant J as $\mathcal{VT}\mathcal{E}(J)$, where $\mathcal{VT}$ is the volume of Euclidean space-time, and $\mathcal{E}(J)$ is the vacuum energy density in the presence of the constant source J . In this formalism, $J(x)$ plays a role analogous to that of a spatially dependent external magnetic field in a ferromagnet, and $\langle \phi(x) \rangle$ is the analog of the local magnetization at x . Spontaneous symmetry breaking is typically determined from the solution of the relation

$$\langle \phi(x) \rangle = -\frac{\delta W}{\delta J(x)}_{J=0}. \quad (16)$$

This is generally not a closed equation for $\langle \phi(x) \rangle$, and approximation solutions from, *e.g.*, perturbation theory must be used. The functional $W[J]$ is the field-theoretic analog of the free energy of a spin system in an external magnetic field. It will be convenient in what follows to work with ϕ as the independent variable rather than J . The effective action $\Gamma[\phi_c]$ is a functional of the classical field $\phi_c(x)$ defined by

$$\phi_c(x) = -\frac{\delta W}{\delta J(x)} \quad (17)$$

for arbitrary $J(x)$. Given that relation between $J(x)$ and $\phi_c(x)$, we define Γ via a functional Legendre transform:

$$\Gamma[\phi_c] = W[J] + \int d^{d+1}x J(x) \phi_c(x) \quad (18)$$

which satisfies

$$\frac{\delta \Gamma}{\delta \phi(x)} = J(x). \quad (19)$$

This construction is precisely analogous to exchanging the role of independent variable between the external field and the magnetization in a ferromagnet. For constant fields, $W[J]$ is the free energy of the system with constant J , and $\Gamma[\phi_c]$ is the free energy of the system with the conjugate variable ϕ_c held constant. For the case of a single scalar field, the effective potential is given at one loop as

$$\Gamma[\phi_c] = S[\phi_c] + \frac{\hbar}{2} \text{Tr} [\log (-\nabla^2 + V''(\phi_c))] \quad (20)$$

Γ also has a graphical interpretation as the generator of one-particle-irreducible (1PI) Feynman graphs.

In many applications, it is convenient to consider the effective potential $V_{eff}(\phi_c)$. It can be obtained from a derivative expansion of $\Gamma[\phi_c]$ for slowly-varying fields:

$$\Gamma[\phi_c] = \int d^{d+1}x \left[V_{eff}(\phi_c) + \frac{1}{2} Z(\phi_c) (\partial_\mu \phi_c)^2 + \dots \right] \quad (21)$$

where quartic-derivative terms and higher are indicated by ellipses. The effective potential should be understood at zero temperature as the vacuum energy density of the system given that $\langle \phi(x) \rangle = \phi_c$. We have for constant fields

$$\frac{\partial V_{eff}}{\partial \phi_c} = J \quad (22)$$

and $V_{eff}(\phi_c)$ is related to $\mathcal{E}(J)$ by a Legendre transform

$$V_{eff}(\phi_c) = \mathcal{E}(J) + J\phi_c. \quad (23)$$

It is important to note that this construction is ambiguous when there are multiple solutions for ϕ_c . In general, $V_{eff}(\phi_c)$ is given by the convex hull of the differential construction, a generalization of Maxwell's equal-area construction.

V_{eff} is an extension of the classical potential V including quantum effects. From the point of view of perturbation theory, $V_{eff} = V + O(\hbar)$, and the $O(\hbar)$ correction term has

$$V_{eff}(\phi_c) = V(\phi_c) + \frac{\hbar}{2} \int \frac{d^4k}{(2\pi)^4} \log(k^2 + V''(\phi_c)) \quad (24)$$

Note that this expression requires renormalization. It is easy to show that this expression may also be written as

$$V_{eff}(\phi_c) = V(\phi_c) + \hbar \int \frac{d^3k}{(2\pi)^3} \frac{1}{2} \sqrt{k^2 + V''(\phi_c)} \quad (25)$$

which clearly exposes the one-loop correction as a sum over the zero-point energies of all the field modes. The background field method offers a very powerful technique for evaluating $\Gamma[\phi]$.

E. Field theory at finite temperature

In quantum statistical mechanics, the partition function is given by $Z = \text{Tr}[\exp(-\beta H)]$ where β is the inverse of the temperature T . The propagation in Euclidean time is over a finite extent, given by β , and the trace can be implemented by requiring periodic boundary conditions on the fields. The partition function is just the generating function of the field theory

$$Z = \int [d\phi] e^{-S} \quad (26)$$

but the functional integral is now over all field configurations satisfying $\phi(\vec{x}, 0) = \phi(\vec{x}, \beta)$. Thus the effects of non-zero temperature are included by changing the geometry of the system from R^{d+1} to $R^d \times S^1$ with periodic boundary conditions for bosons; fermions require antiperiodic boundary conditions.

The generalization of the one-loop effective potential to finite temperature is illuminating. Due to the periodic boundary conditions in the Euclidean time direction, the integral over k_{d+1} is replaced by a sum over Matsubara frequencies

$$V_{eff}(\phi_c) = V(\phi_c) + \frac{\hbar}{2\beta} \sum_{n \in \mathbb{Z}} \int \frac{d^3 k}{(2\pi)^3} \log \left(\left(\frac{2\pi n}{\beta} \right)^2 + \vec{k}^2 + V''(\phi_c) \right) \quad (27)$$

which can be converted into the form

$$V_{eff}(\phi_c) = V(\phi_c) + \hbar \int \frac{d^3 k}{(2\pi)^3} \left[\frac{1}{2} \sqrt{\vec{k}^2 + V''(\phi_c)} + \log \left(1 - e^{-\beta \sqrt{\vec{k}^2 + V''(\phi_c)}} \right) \right] \quad (28)$$

The additional term is instantly recognizable as $-p$, where p is the pressure of a relativistic Bose gas with mass $\sqrt{V''(\phi_c)}$. This is of course consistent with the identification of $V_{eff}(\phi_c)$ with the free energy of the system, when evaluated at its global minimum.

F. Symmetry restoration at high temperatures

Generally speaking, broken symmetries are associated with low temperatures, as seen, for example, in spin systems such as the Ising model. This can be explicitly seen in scalar field theories at finite temperature using the effective potential. The finite temperature part of the one-loop effective potential has a natural expansion for high temperatures. A more general form for such expressions will be given in section III C, so for now we simply note

that that for a single real scalar field the first two terms in a high-temperature expansion are given by [9, 10]

$$V_{1T} \simeq -\frac{\pi^2 T^4}{90} + \frac{1}{24} T^2 V''(\phi_c). \quad (29)$$

The first term is the usual blackbody term with no effect on the value of ϕ_c . However, the sub-leading T^2 term acts to restore symmetry. If the potential $V(\phi)$ has the form

$$V(\phi) = -\frac{1}{2}\mu^2\phi^2 + \frac{1}{4!}\lambda\phi^4, \quad (30)$$

then $\mu^2 > 0$ will lead to spontaneous symmetry breaking at zero temperature. However V_{1T} supplies a temperature-dependent mass term such that the potential for ϕ becomes

$$V(\phi) + V_{1T}(\phi) \simeq -\frac{\pi^2 T^4}{90} + \frac{1}{2} \left(\frac{\lambda}{24} T^2 - \mu^2 \right) \phi^2 + \frac{1}{4!} \lambda \phi^4 \quad (31)$$

and the symmetry $\phi \rightarrow -\phi$ will be restored at temperatures above $T_c \simeq \sqrt{24/\lambda}\mu$. A notable exception to the general rule of symmetry restoration at high temperature is the deconfinement transition in pure gauge theories, where the broken phase lies above the unbroken phase in temperature. This will be explain in detail in Section III B.

The temperature dependence of the effective potential is very similar to the general form of the free energy as a function of order parameter postulated by Landau theory. A slightly different construction, often referred to as dimensional reduction, also leads to a free energy of Landau's form. In dimensional reduction, the field modes with non-zero Matsubara frequencies are integrated out, so that only modes which are constant in Euclidean time remain. This effectively reduces a $(d+1)$ -dimensional theory to a d -dimensional theory with temperature-dependent parameters in the action.

G. Spatially modulated phases

A further complication in the determination of the phase diagram is the possible existence of spatially modulated phases. In the most familiar cases, symmetry breaking occurs when some order parameter $\phi(x)$ has a constant, non-zero expectation value independent of x . However, there are phases in nature where order parameters are not translationally and/or rotationally invariant. The most familiar example is the formation of crystalline solid phases in ordinary materials, where the order parameter is the density. In a liquid or vapor phase, the density is uniform, but in a crystalline phase it is periodic. Magnetic systems can also

show spatially modulated phases, as exhibited by simple models such as the anisotropic next-nearest-neighbor Ising (ANNNI) model [11] and the chiral potts model [12, 13]. In field theories, spatially modulated phases are generally associated with systems at finite density, and the sign problem is intimately involved in this behavior [14, 15]. In QCD, some color superconducting phases are crystalline in nature [16]; these are analogs of the Larkin-Ovchinnikov-Fulde-Ferrell (LOFF) phases of ordinary superconductors [17, 18].

There is a simple approach to spatially modulated phases due to Lifshitz [3]. As a simple model, consider a Landau-Ginsburg model with a free energy density of the form

$$f(\phi) = \frac{a}{2} (\nabla\phi)^2 + \frac{b}{2} (\nabla^2\phi)^2 + \frac{r}{2}\phi^2 + \frac{\lambda}{4}\phi^4 \quad (32)$$

where b and λ must be positive but a and r can be of either sign. Taking the case $a < 0$ and $r > 0$, it is easy to see that the homogeneous equilibrium state will be unstable to spatially-varying perturbations of wavenumber k for any values of k satisfying $bk^4 + ak^2 + r < 0$. This model has three phases: an unbroken phase, a broken phase, and a spatially modulated phase. The point $a = 0, r = 0$ is the Lifshitz point, where the three phases coexist.

H. Non-equilibrium behavior: nucleation, spinodal decomposition and relaxation

The perturbative effective potential can yield information about non-equilibrium behavior. Recall that the perturbative effective potential does not satisfy the convexity properties required of V_{eff} by its Legendre transform construction. However, the non-convex region of V_{eff} contains information about non-equilibrium behavior. Metastable states can be understood as local minima of the effective potential. Consider a field theory with a double well potential of the form $V(\phi) = \lambda(\phi^2 - v^2)^2$. The presence of a small additional linear coupling $-J\phi$ will bias the system towards one of the two minima at $\pm v$. This leads to a discontinuity in the behavior of ϕ_c as a function of J . However, the perturbative effective potential shows the persistence of metastable states across $J = 0$. The intuitive picture is simple in the thin wall approximation, which has roots in classical nucleation theory. Consider an initial, homogeneous field configuration where $\phi = \phi_{local}$, the value of the local minimum of V_{eff} . Coherent fluctuations in space will produce small droplets, also called bubbles, of ϕ_{global} , the global minimum. The difference between the free energy densities $\Delta f = f(\phi_{local}) - f(\phi_{global})$ drives the expansion of such bubbles. However, there is also a surface tension σ residing in

the interface between the two phases, which opposes the growth of bubbles. In the thin-wall approximation, the free energy of a three-dimensional droplet of radius R is approximately

$$F(R) = -\Delta f \cdot \frac{4\pi}{3} R^3 + \sigma \cdot 4\pi R^2 \quad (33)$$

where σ is the interfacial surface tension at equilibrium. The surface tension may be calculated from the one-dimensional kink solution that interpolates between the two phases. A droplet which is smaller than the critical size R_c will shrink due to surface tension effects, while a droplet with $R > R_c$ will expand to convert all space to the equilibrium phase. The decay rate of metastable states can be calculated within the functional integral formalism using instanton techniques [19–21].

$$\Gamma = A e^{-S_b} \quad (34)$$

where S_b is the action of the so-called bounce solution. Within the thin-wall approximation, it is equal to $F(R_c)$. The pre-factor A is given in terms of a functional determinant representing fluctuations around the bounce solution.

Within the saddle-point approximation considered here, there is a clear point at which the metastable phase ceases to exist, *i.e.*, where the metastable states cease to exist as local minima. The subspace of parameter space where this occurs is referred to as the spinodal, which may be a point, line, *et cetera*. The spinodal can be located by the condition $V'_{eff}(\phi_{spinodal}) = V''_{eff}(\phi_{spinodal}) = 0$, and more generally by the vanishing of a mass associated with a metastable state. In physical systems, fluctuations become important near the spinodal point, and the distinction between metastable versus unstable states becomes blurred.

Unstable states are of interest in their own right, and something about their behavior can be learned using the effective action. Consider a ϕ^4 field theory in d spatial dimensions at finite temperature. We apply dimensional reduction so we have a three-dimensional field theory with temperature-dependent coefficients in the potential. Suppose the system begins in a high-temperature equilibrium state with unbroken symmetry so $\phi = 0$, but has a stable broken symmetry phase at low temperatures. A rapid lowering of the temperature below the critical point, known as a rapid quench, will place the system in a state of unstable equilibrium, with $V''_{eff}(\phi = 0) < 0$. As in the case of a metastable state, such an unstable

state will lead to a free energy

$$f(\phi = 0) = V_{eff}(\phi = 0) = V(\phi = 0) + \frac{1}{2} \int \frac{d^d k}{(2\pi)^d} \log(\vec{k}^2 + V''(\phi = 0)) \quad (35)$$

Because $V''_{eff}(\phi = 0) < 0$, the modes near $\vec{k} = 0$ in the functional determinant give rise to an imaginary component of the free energy. This imaginary part can be associated with the decay rate of the unstable phase in a manner similar to the decay of metastable states [22]. In the early stages of equilibration, the unstable modes in the region $\vec{k}^2 + V''(\phi = 0) < 0$ lead to exponentially growth of long-wavelength oscillations in the order parameter. This process is known as spinodal decomposition. When the order parameter is not conserved, *e.g.* in the case of magnetization, the fastest growth occurs around $\vec{k} = 0$; however, in the case of a conserved order parameters, such as a charge density, the fastest growth occurs at a non-zero value of $|\vec{k}|$ [3].

In addition to nucleation and spinodal decomposition, there is a third equilibration process, one which is always present in non-equilibrium situations. Relaxation is the process whereby a system returns to its equilibrium value after a small perturbation away from equilibrium; the term may also be applied to individual field modes. An introductory discussion of relaxation requires a treatment of damping, *i.e.* frictional or dissipative terms in the effective action. All three processes have been extensively discussed in the context of the early universe [23, 24], particularly with respect to inflation.

III. PHASES OF FINITE TEMPERATURE GAUGE THEORIES

Gauge theories with $T \neq 0$ (“finite temperature”) have a rich phase structure which has been extensively explored using a combination of analytic methods and lattice simulations. Non-Abelian gauge theories have global symmetries and associated order parameters which are analogous to magnetization in spin systems, and much of the modern formalism of critical phenomena is directly applicable. The symmetries and order parameters associated with quark confinement and chiral symmetry breaking are of particular interest as principal determinants of gauge theory phase structure.

A. Confinement

The most striking feature of QCD is confinement: the force between widely separated quark-antiquark pairs is a constant σ , known as the string tension. This constant force implies a potential energy between a quark-antiquark pair that grows as σr for large distances r . The numerical value of σ , determined from phenomenology, is approximately $0.18 \text{ GeV}^2 \approx 0.9 \text{ GeV}/\text{fm}$. The string tension can be extracted from the asymptotic behavior of the quark-antiquark potential. In turn, the static potential between two heavy quarks can be determined in a gauge-invariant way from the expectation value of the Wilson loop. This is a non-local operator associated with a closed curve $\mathcal{C}$ in space-time parametrized as $x_\mu(\tau)$ where τ can be taken to be the unit interval and $x_\mu(1) = x_\mu(0)$. The Wilson loop is defined as

$$W[\mathcal{C}] = \mathcal{P} \exp \left[i \oint_{\mathcal{C}} dx_\mu A_\mu(x) \right] \quad (36)$$

where $\mathcal{P}$ indicates path-ordering of the gauge fields $A_\mu(x)$ along the path $\mathcal{C}$ [25]. Taking $W[\mathcal{C}]$ to be an element of the abstract gauge group G , the basic observables are $\text{Tr}_R W[\mathcal{C}]$, where the trace is taken over an irreducible representation R of G . $\text{Tr}_R W[\mathcal{C}]$ has a physical interpretation as the non-Abelian phase factor associated with a heavy particle in the representation R moving adiabatically around the closed loop $\mathcal{C}$. Typically, we are interested in rectangular Wilson loops of width sides L and T . The loop can be associated with a process in which a particle-antiparticle pair are created at one time, move to a separation L , propagate forward in time for an interval T , and then annihilate. For $T \gg L$, we have

$$\langle \text{Tr}_R W[\mathcal{C}] \rangle \simeq \exp[-V_R(L)T] \quad (37)$$

where $V_R(L)$ is the heavy quark-antiquark potential for particles in the representation R . For large distances, the potential can grow no faster than linearly with L , and that linear growth defines the string tension σ_R for the representation R :

$$V(L) \rightarrow \sigma_R L + \mathcal{O}(1) \quad (38)$$

as $L \rightarrow \infty$. The $\mathcal{O}(1)$ correction term represents a perimeter-law contribution, which must be renormalized; the effect is closely related to the mass renormalization of a heavy particle interacting with the gauge field.

Beyond the perimeter law contribution, there are corrections which are $\mathcal{O}(1/L)$ and higher that are of physical interest. Using an effective string description, Luscher showed

that the $\mathcal{O}(1/L)$ has a universal coefficient that depends only on the dimensionality of space-time [26]; this prediction was confirmed in lattice simulations [27]. Subsequent analytical work has explored the effects of higher-order terms [28–30]. There have also been results on the broadening of the flux tube as a function of L [31, 32]. For a lattice-oriented review of string effects in gauge theories, see the review by Pepe [33].

The statement that quarks are confined is the statement that the string tension σ_F of the fundamental representation of $SU(3)$ gauge theory is non-zero. However, each representation has its own string tension. In lattice simulations of $SU(3)$, the fundamental string tension is the smallest, and representations of larger dimension generally have a larger string tension. A characteristic feature of gauge theories in four dimensions is dimensional transmutation: the classical action is scale-invariant, but the introduction of a mass scale μ is necessary in order to define the running coupling constant $g(\mu)$. It follows directly from the renormalization group that any physical mass in four-dimensional pure gauge theories must be proportional to a renormalization group invariant mass, usually written simply as Λ . This implies that $\sigma_R = c_R \Lambda^2$, where c_R is a pure number. Put another way, the ratio σ_R/σ_F is a pure number characterizing the representation R . In (1+1)-dimensional gauge theories, the string tension σ_R is proportional to the quadratic Casimir invariant, leading to a behavior known as Casimir scaling. It is generally believed that a satisfactory description of confinement in four dimensions would include an understanding of the string-tension scaling law.

There are two subtleties associated with the string tension. The first subtlety is that string tensions are generally determined in so-called pure gauge theories, where only the gauge fields are dynamical; quarks and other particles exist only as static classical charges. This is done because of string breaking, also known as charge screening. Consider a theory with dynamical particles of mass m in the fundamental representation F of the gauge group. Then energy obtained by separating a pair of static sources in the fundamental representation is $\sigma_F L$; when that energy becomes on the order of $2m$, it will become energetically favorable to produce a pair of dynamical particles at a cost in energy of $2m$, and no string tension will be seen for $L \gtrsim 2m/\sigma_F$.

The second subtlety is related: in $SU(N)$ gauge theories, a state with $N + 1$ fermions in the fundamental representation, *e.g.* quarks in $SU(3)$, will form a baryon, a neutral state of N fermions, plus one additional fermion. Thus the string tension between $N + 1$ fermions and $N + 1$ antifermions observed at large distances should be the same string tension observed

between one fermion and one antifermion. Thus for purposes of confinement, the relevant “charge” is an integer k which can be taken between 1 and N . Furthermore, the string tension is the same for states of N -ality k and $N - k$, so that there are $\lfloor \frac{N}{2} \rfloor$ independent string tensions. This behavior is associated with the $Z(N)$ symmetry underlying confinement in $SU(N)$, discussed below.

It is generally believed that there is some simple regular behavior described by string tension scaling laws [34]. For example, in $(1 + 1)$ -dimensions, Casimir scaling gives

$$\frac{\sigma_k}{\sigma_1} = \frac{k(N - k)}{N - 1} \quad (39)$$

This behavior is also found in the strong-coupling limit of lattice gauge theories in all dimensions, and is consistent with the behavior seen in lattice simulations. An alternative to Casimir scaling is sine-law scaling

$$\frac{\sigma_k}{\sigma_1} = \frac{\sin\left(\frac{\pi k}{N}\right)}{\sin\left(\frac{\pi}{N}\right)} \quad (40)$$

a behavior predicted in certain supersymmetric gauge theories [35], in MQCD [36]. This behavior is also roughly consistent with lattice results. Note that both formulas give $\sigma_N = 0$ and are invariant under $k \leftrightarrow N - k$. In the limit $N \rightarrow \infty$ at fixed k , both formulas give $\sigma_k \rightarrow k\sigma_1$. Corrections to the large- N limit start at $1/N$ for Casimir scaling and $1/N^2$ for sine-law scaling, and it has been argued that the large- N limit requires $1/N^2$ corrections [37]. However, lattice simulations of $SU(N)$ gauge theories in $(2 + 1)$ dimensions indicate that $1/N$ corrections are strongly indicated [38], and a careful analysis shows that Casimir scaling is compatible with the large- N limit [39].

B. Pure gauge theories at finite temperature

If one or more directions in space-time are compact, the string tension may also be measured using the Polyakov loop P , also known as the Wilson line. The Polyakov loop is essentially a Wilson loop that uses a compact direction in space-time to close the curve using a topologically non-trivial path in space time, as shown in figure 2. The typical use for the Polyakov loop is for gauge theories at finite temperature, where space-time is $R^3 \times S^1$. The partition function being given by $Z = \text{Tr} [e^{-\beta H}]$, the circumference of S^1 is given by




Figure 2: The Polyakov loop is associated with the worldline of a heavy particle.

the inverse temperature $\beta = 1/T$. In this case, we write

$$P(\vec{x}) = \mathcal{P} \exp \left[i \int_0^\beta dx_4 A_4(x) \right] \quad (41)$$

and string tensions may be determined from a two-point function

$$\langle Tr_R P(\vec{x}) Tr_R P^\dagger(\vec{y}) \rangle \sim e^{-\beta \sigma_R |\vec{x} - \vec{y}|} \quad (42)$$

a behavior that assumes that the one-point function $\langle Tr_R P(\vec{x}) \rangle = 0$. This behavior is shown in figure 3. The Polyakov loop one-point function $\langle Tr_R P(\vec{x}) \rangle$ can be interpreted as a Boltzmann factor $\exp(-\beta F_R)$, where F_R is the free energy required to add a static particle in the representation R to the system. Of course, $\langle Tr_R P(\vec{x}) \rangle = 0$ implies that $F_R = \infty$, which is thus a fundamental criterion determining whether particles in the representation R are confined. Note that the introduction of a compact direction breaks the four-dimensional symmetry of the theory, and the string tension measured by Polyakov loops is not the same as the string tension measured by Wilson loops lying in non-compact planes. In the case of finite temperature, it is natural to use the terminology electric and magnetic string tension, respectively. In the limit where the compactification radius becomes large, *e.g.* $\beta \rightarrow \infty$, the two string tensions must coincide.

1. Center symmetry

One of the most important concepts in our understanding of confinement is the role of center symmetry. The center of a Lie group is the set of all elements that commute with every other element. For $SU(N)$, this is $Z(N)$. Although the $Z(N)$ symmetry of $SU(N)$




Figure 3: The Polyakov loop two-point function determines the electric string tension.

gauge theories can be understood from the continuum theory, it is easier to understand from a lattice point of view. A lattice gauge theory associates link variable $U_\mu(x)$ with each lattice site x and direction μ . The link variable is considered to be the path-ordered exponential of the gauge field from x to $x + \hat{\mu}$: $U_\mu(x) = \exp[iaA_\mu(x)]$. Consider a center symmetry transformation on all the links in a given direction on a fixed hyperplane perpendicular to the direction. The standard example from $SU(N)$ gauge theories at finite temperature is $U_4(\vec{x}, t) \rightarrow zU_4(\vec{x}, t)$ for all $\vec{x}$ and fixed t , with $z \in Z(N)$. Because lattice actions such as the Wilson action consist of sums of small Wilson loops, they are invariant under this global symmetry. However, the Polyakov loop transforms as $P(\vec{x}) \rightarrow zP(\vec{x})$, and more generally

$$Tr_R P(\vec{x}) \rightarrow z^{k_R} Tr_R P(\vec{x}) \quad (43)$$

where k_R is an integer in the set $\{0, 1, \dots, N-1\}$ and is known as the N -ality of the representation R . If $k_R \neq 0$, then unbroken global $Z(N)$ symmetry implies $\langle Tr_R P(\vec{x}) \rangle = 0$. Thus global $Z(N)$ symmetry defines the confining phase of a gauge theory. For pure gauge theories at non-zero temperature, the deconfinement phase transition is associated with the loss of $Z(N)$ symmetry at the critical point T_d . Below that point $\langle Tr_F P(\vec{x}) \rangle = 0$ but above T_d , $\langle Tr_R P(\vec{x}) \rangle \neq 0$.

Notice that the case of zero N -ality representations is special within this framework: there is no requirement from $Z(N)$ symmetry that these representations are confined. This includes the adjoint representation, the representation of the gauge particles. However, lattice simulation indicate that $\langle Tr_R P(\vec{x}) \rangle$ is very small for these representations in the confined phase. Although screening by gauge particles must dominate at large distances, these zero N -ality representations have well-defined string tensions at intermediate distances

scales, *e.g.*, on the order of a few fermi for $SU(3)$, behaving in a manner very similar to representations with non-zero N -ality [40, 41].

2. The nature of the confined phase and partially confined phases

The requirements for confinement are simple: $Z(N)$ symmetry and a mass gap. However, gauge theories with $SU(N)$ symmetry are not simply $Z(N)$ systems. In models where the order parameter is an element of $Z(N)$, the order parameter must average to zero in the disordered phase, and the equilibrium disordered state must consist of many domains, each with its own $Z(N)$ orientation. This need not be the case with $SU(N)$ systems. There is a unique set of Polyakov loop eigenvalues that are $Z(N)$ symmetric, and can form the basis for a semiclassical understanding of confinement.

Although $Tr_F P$ is the order parameter for confinement of particles in the fundamental representation, it does not by itself characterize the confined phase, and the most general form of the free energy does not depend on P solely through $Tr_F P$ and $Tr_F P^\dagger$ [42]. The high-temperature effective potential $V_{eff}(P)$ illustrates this point. It is not a function solely of $Tr_F P$ and its conjugate, and cannot be written as an infinite series in $Tr_F P$ and $Tr_F P^\dagger$. A simple example will illustrate this point. Consider two diagonal matrices lying in $SU(4)$, defined by

$$P_1 = \begin{pmatrix} e^{i\pi/4} & & & \\ & e^{i3\pi/4} & & \\ & & e^{i5\pi/4} & \\ & & & e^{i7\pi/4} \end{pmatrix} \quad (44)$$

and

$$P_2 = \begin{pmatrix} e^{i\pi/2} & & & \\ & e^{i\pi/2} & & \\ & & e^{-i\pi/2} & \\ & & & e^{-i\pi/2} \end{pmatrix} \quad (45)$$

Both P_1 and P_2 have zero trace in the fundamental representation, yet the traces of their square are different: $Tr_F P_1^2 = 0$ and $Tr_F P_2^2 = -4$, establishing that $Tr_F P^2$ cannot be a function solely of $Tr_F P$.

At first sight, this may seem to contradict two standard results: a) the characters form a complete, in fact orthogonal, basis on class functions; b) all characters may be obtained from the fundamental representation by repeated multiplication and the application of

$$\chi_a(P)\chi_b(P) = \sum_c n(a, b; c) \chi_c(P) \quad (46)$$

where all n 's are non-negative integers. Taken together, these results might suggest that all group characters are polynomials in $Tr_F U$ and its complex conjugate. Consider, however, the product representation $N \otimes N$. It is reducible into $N(N+1)/2 \oplus N(N-1)/2$, which are symmetric and antisymmetric representations, respectively. Their characters are respectively

$$\begin{aligned} \chi_S(P) &= \frac{1}{2} [(Tr_F P)^2 + Tr_F(P^2)] \\ \chi_A(P) &= \frac{1}{2} [(Tr_F P)^2 - Tr_F(P^2)]. \end{aligned} \quad (47)$$

Note that the sum $\chi_S + \chi_A$ is a polynomial in $Tr_F P$, but χ_S and χ_A are in general not. For example, the restriction on the eigenvalues of P imposed by $\det(P) = 1$ allows us to prove that for $SU(3)$

$$\frac{1}{2} [(Tr_F P)^2 - Tr_F(P^2)] = Tr_F(P^\dagger) \quad (48)$$

in accord with the $SU(3)$ result $3 \otimes 3 = 6 \oplus \bar{3}$. In this case, it is true that $Tr_F(P^2)$ can be written as a polynomial in $Tr_F P$ and $Tr_F P^\dagger$. However, for $SU(4)$, unitarity of P gives instead

$$\frac{1}{2} [(Tr_F P)^2 - Tr_F(P^2)] = \frac{1}{2} [(Tr_F P)^2 - Tr_F(P^2)]^* \quad (49)$$

which shows that the 6 representation of $SU(4)$ is real. $SU(N)$ group characters can be written as polynomials in $Tr_F(P)$ and its complex conjugate only for $SU(2)$ and $SU(3)$.

An alternative statement is that the Polyakov loop in the fundamental representation, $Tr_F U$, is not sufficient to determine the eigenvalues of P , beginning with the case of $SU(4)$. Let us label the eigenvalues for the Polyakov loop as z_i , with $i = 1$ to N . For $SU(3)$, the characteristic polynomial for the Polyakov loop is

$$\prod_{i=1}^3 (z - z_i) = z^3 - (z_1 + z_2 + z_3) z^2 + (z_1 z_2 + z_2 z_3 + z_3 z_1) z - z_1 z_2 z_3. \quad (50)$$

Since the determinant of a special unitary matrix is 1, we have $z_1 z_2 z_3 = 1$, and the characteristic polynomial is

$$z^3 - z^2 Tr_F P + z Tr_F P^\dagger - 1 \quad (51)$$

so for $SU(3)$, knowledge of $Tr_F P$ determines all eigenvalues, and the free energy can be written as a function of $Tr_F P$ alone. For $SU(4)$, similar considerations allow the characteristic polynomial to be written as

$$z^4 - z^3 Tr_F P + z^2 \frac{1}{2} [(Tr_F P)^2 - Tr_F (P^2)] - z Tr_F P^+ + 1 \quad (52)$$

and knowledge of $Tr_F P$ must be supplemented by the value of $Tr_F (P^2)$. As N increases, more information must be supplied to reconstruct the eigenvalues.

The characteristic equation for Polyakov loops in $SU(N)$ may be written as

$$\det [P - \lambda I] = \sum_{j=0}^N (-\lambda)^j \chi_j(P) \quad (53)$$

where χ_j is the character for the representation formed from the antisymmetrized product of the fundamental j times; the corresponding Young tableaux is j vertical boxes. Of course, $\chi_j(P)$ has N -ality j , and $\chi_0(P) = \chi_N(P) = 1$. At the classical level, a Polyakov loop that represents the confining phase must have $\chi_j(P) = 0$ for $1 \leq j \leq N-1$, leading to a characteristic equation $1 + (-\lambda)^N = 0$. This in turn implies the important result that any model in which confinement is associated with a particular field configuration (up to gauge transformations) must have Polyakov loop eigenvalues evenly spaced on the unit circle in the confined phase. This is precisely what must happen in the large- N limit, and is known to occur in analytically tractable lattice models of the large- N limit [43]. This behavior is not possible in simple $Z(N)$ models where symmetric phases always represent some average over different $Z(N)$ domains. However, it should be noted that there are $Z(N)$ spin systems with features similar to models with variables in $SU(N)$. The best-known is the Blume-Emery-Griffiths model, which was originally proposed as a simple model of ${}^3\text{He}$ - ${}^4\text{He}$ mixtures [44]. Essentially, an Ising spin model is extended such that each spin σ can have the value 0 as well as ± 1 . The behavior of such extended $Z(N)$ lattice models is very similar to treatments of $SU(N)$ lattice models with fundamental and adjoint interactions, where the adjoint interaction can be adjusted to give a $Z(N)$ limit [45, 46]; see also [47].

In the confining phase of a gauge theory, $Z(N)$ symmetry is unbroken, and all representations with non-zero N -ality are confined. In the deconfined phase, $Z(N)$ symmetry is completely lost, and particles are no longer confined, independent of their representation. For $N \geq 4$, additional phases are possible where $Z(N)$ is broken down to a non-trivial subgroup [2, 42, 48, 49]. In the case of $Z(4)$, there can be breaking of center symmetry down to

$Z(2)$. In this partially confined phase, states consisting of two fundamental representation fermions are not confined, but single fermions are. In the case of $SU(N)$, $Z(N)$ can break to $Z(k)$, where k is any divisor of N . States with 1 to $k-1$ fermions are confined, but states with k fermions are not. It is often convenient to include the confined and deconfined phases as $k = N$ and $k = 1$, respectively. As we will see in section IV, such partially confined phases have been found in gauge theories on $R^3 \times S^1$, using both lattice simulations and perturbation theory [2]. It should be noted that not all gauge groups have non-trivial centers. The gauge group $G(2)$ provides an interesting example of a gauge theory without a center [50] that has received significant attention [51].

3. Origin of the deconfinement transition

As we have seen in section II, the critical properties of a four-dimensional field theory at finite temperature can often best be understood in terms of an effective three-dimensional theory. From this point of view, the Polyakov loop can be considered to be a scalar order parameter with $Z(N)$ invariance in some effective three-dimensional theory. A complete treatment of the effective potential, including non-perturbative effects, would presumably yield the phase structure, including both the confined and deconfined phases.

Perturbation theory is a reliable indicator of broken center symmetry and thus deconfinement at high temperature, because the running coupling constant $g(T)$ is small if $T \gg \Lambda$. The one-loop effective potential for a pure gauge theory in the background of a static Polyakov loop P can be easily evaluated in a gauge where A_4 is time-independent and diagonal [52, 53]. It is easy to see that $V_{eff}^{1l}(P)$ is given by

$$V_{eff}^{1l}(P) = 2T \text{Tr}_{Adj} \sum_{n \in Z} \int \frac{d^3k}{(2\pi)^3} \log \left[(2\pi nT - A_4)^2 + \vec{k}^2 \right] \quad (54)$$

where the factor of 2 represents the two helicity states of each mode. Note that there is no classical contribution. Discarding the zero-point energy term, we obtain the one-loop finite-temperature effective potential for gauge bosons

$$V_{eff}^{1l}(P) = 2T \text{Tr}_{Adj} \int \frac{d^3k}{(2\pi)^3} \log \left[1 - P \exp \left(- \left| \vec{k} \right| / T \right) \right] \quad (55)$$

which is the free energy density of the gauge bosons in the background of the Polyakov loop P .

The logarithm in this expression for $V_{eff}^{1l}(P)$ can be expanded, leading to an interpretation of $V_{eff}^{1l}(P)$ as a sum of contributions from gluon worldlines wrapping around the compact direction an arbitrary number of times. Explicitly, we have the expression

$$V_{eff}^{1l}(P) = -\frac{2}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^4} Tr_{Adj} P^n. \quad (56)$$

From this form, it is easy to see that $V_{eff}^{1l}(P)$ is minimized when all the moments $Tr_A P^n$ are maximized. This occurs when $P \in Z(N)$, which gives $Tr_A P^n = N^2 - 1$. This indicates that the one-loop gluon effective potential favors the deconfined phase. The pressure p is the negative of the free energy density at the minimum,

$$p(T) = 2(N^2 - 1) \frac{\pi^2 T^4}{90}, \quad (57)$$

which is exactly p for a blackbody with $2(N^2 - 1)$ degrees of freedom.

In the gauge where A_4 is diagonal and time-independent, we can parametrize A_4 in the fundamental representation of $SU(N)$ as a diagonal, traceless $N \times N$ matrix

$$(A_4)_{jk} = T\theta_j \delta_{jk} \quad (58)$$

so that

$$P_{jk} = e^{i\theta_j} \delta_{jk} \quad (59)$$

with $\sum_{j=1}^N \theta_j = 0$. Using the decomposition $F \otimes \bar{F} = 1 \oplus Adj$ and the corresponding decomposition

$$Tr_F P Tr_P P^+ = 1 + Tr_{Adj} P, \quad (60)$$

one realizes that the N^2 eigenvalues of the product representation $F \otimes \bar{F}$ have the form $\exp[i\Delta\theta_{jk}]$, where we define $\Delta\theta_{jk} \equiv \theta_j - \theta_k$. The effective potential is given by

$$V_{eff}^{1l}(P) = -\frac{2}{\pi^2} \sum_{j,k=1}^N \left(1 - \frac{1}{N} \delta_{jk}\right) \sum_{n=1}^{\infty} \frac{1}{n^4} \exp[in\Delta\theta_{jk}]. \quad (61)$$

The infinite sum over n may be carried out explicitly in terms of the fourth Bernoulli polynomial. For our purposes, a convenient explicit form is

$$V_{eff}^{1l}(P) = -T^4 \sum_{j,k=1}^N \left(1 - \frac{1}{N} \delta_{jk}\right) \left[\frac{\pi^2}{45} - \frac{1}{24\pi^2} |\Delta\theta_{jk}|_{2\pi}^2 (|\Delta\theta_{jk}|_{2\pi} - 2\pi)^2 \right] \quad (62)$$

where $|\Delta\theta_{jk}|_{2\pi}$ lies in the interval between 0 and 2π .

In many systems, broken symmetry phases are found at low temperatures and symmetry is restored at high temperatures. The phase structure of gauge theories as a function of temperature is unusual because the broken-symmetry phase is the high-temperature phase. A lattice construction of the effective action for Polyakov loops, valid for strong-coupling, is instructive [54–57]. The spatial link variables may be integrated out exactly if spatial plaquette interactions are neglected. Each spatial link variable then appears only in two adjacent temporal plaquettes, and may be integrated out exactly using the same techniques that are used in the Migdal-Kadanoff real-space renormalization group [43, 55]. The resulting effective action has the form

$$S_{eff} = \sum_{\langle jk \rangle} K \left[\text{Tr}_F P_j \text{Tr}_F P_k^\dagger + \text{Tr}_F P_k \text{Tr}_F P_j^\dagger \right] \quad (63)$$

where K is a function of the lattice gauge coupling g^2 and the extent of the lattice in the Euclidean time direction n_t , which is related to the temperature by $n_t a = 1/T$. In the strong-coupling limit of the underlying gauge theory, the explicit form for K is $K \simeq (1/g^2 N)^{n_t}$ to leading order. In the weak-coupling limit, a Migdal-Kadanoff bond-moving argument gives $K \simeq 2N/g^2 n_t$. This effective action represents a $Z(N)$ -invariant nearest-neighbor interaction of a spin system where the Polyakov loops are the spins. It depends only on gauge-invariant quantities. Standard expansion techniques show that the $Z(N)$ symmetry is unbroken for small K , and broken for K large. This model explains why the high-temperature phase of gauge theories is the symmetry-breaking phase: the relation between K and the underlying gauge theory parameters is such that K is small at low temperatures, and large at high temperatures, exactly the reverse of a classical spin system where the coupling is proportional to T^{-1} . For small values of n_t , the deconfinement transition can be easily extracted, but the phase transition is in the strong-coupling region and far from the continuum limit. A systematic treatment of strong-coupling corrections has recently been shown to yield values for the critical lattice couplings $\beta_c \equiv 2N/g^2$ for $SU(2)$ and $SU(3)$ that are within a few percent of simulation results for $4 \leq N_t \leq 16$ [58]. For Abelian lattice gauge theories in $(2+1)$ -dimensions, a duality transformation maps the high-temperature symmetry-breaking phase of the gauge theory into the high-temperature unbroken phase of the dual spin system, and the low-temperature symmetry-breaking phase of the spin system into the unbroken phase of the gauge theory, consistent with the general behavior seen using strong-coupling arguments.

4. Universality

The lattice construction of the Polyakov loop effective action is a concrete realization of Svetitsky-Yaffe universality [59], which states that a second-order deconfinement transition in a $(d + 1)$ -dimensional gauge theory is in the universality class of classical spin systems in d dimensions with the same global symmetry. Lattice simulations indicate that all pure $SU(N)$ gauge theories have a deconfining phase transition at some temperature T_d , above which center symmetry is broken. In accordance with predictions based on universality, the deconfinement transition for an $SU(2)$ gauge theory in $3 + 1$ dimensions has been well-established as being in the universality class of the three-dimensional Ising model, exhibiting a second-order transition at T_d . The deconfinement transition for $SU(3)$ in $3 + 1$ dimensions is first-order. This is consistent with Landau-Ginsburg predictions for a system with a $Z(3)$ symmetry. The transitions for $N > 3$ appear to be first-order in $3 + 1$ dimensions as well, with a smooth limit as N goes to infinity [60, 61], so the direct applicability of the universality argument in $3 + 1$ dimensions is more limited than one might have expected.

5. Equation of state

Lattice simulations can be used to determine both the pressure $p(T)$ and the internal energy $\epsilon(T)$, along with the deconfinement transition temperature T_d and related properties of the transition; see *e.g.* [62] and references therein. For pure gauge theories below T_d , both p and ϵ are extremely small. This is expected because the mass of the lightest color singlet state, the scalar glueball, is substantially larger than T_d . The pressure must be continuous at T_d , but the first-order character of the deconfinement transition for $N \geq 3$ leads to a latent heat, indicated by a discontinuity in ϵ . The thermodynamics quantities p and ϵ as well as the order parameter $\langle Tr_F P \rangle$, show a rapid rise in the interval from T_d to roughly $3T_d$. Both p and ϵ show a monotonic but slow rise towards their blackbody values at higher temperatures.

The pressure $p(T)$ and related thermodynamic quantities can be calculated in perturbation theory in terms of $g^2(T)$, the running coupling at temperature T . However, there is a barrier to perturbative calculations at $\mathcal{O}(g^6)$, due to infrared divergences in the magnetic sector [63]. The best that can be done in perturbation theory is an expression for $p(T)$

that is valid up to $\mathcal{O}(g^6 \log g)$ [64]. Comparison of different order of perturbation theory up to $\mathcal{O}(g^6 \log g)$ shows a reasonable convergence only at temperatures greater than about $10\Lambda_{\overline{MS}}$.

6. Phenomenology of the deconfinement transition

The availability of high-quality lattice data for the $SU(3)$ pressure [65–68] has led to a variety of attempts to model it phenomenologically. Deconfinement can be characterized broadly as a change in the number of degrees of freedom as the temperature is raised, and quark and gluon degrees of freedom manifest in thermodynamic behavior. The key to building a successful model is incorporating a mechanism for characterizing the change in degrees of freedom with temperature. Phenomenological models built around minimizing the free energy as a function of the order parameter are appealing and simple [42, 69, 70].

One simple model [42, 71] follows from noting that the one-loop free energy for massive particles in a Polyakov loop background lead to expansions in which the first two terms have simple forms when $M/T \ll 1$. Thus we arrive at a potential of the form

$$V_G(P) = -\frac{2T^4}{\pi^2} \sum_{n=1}^{\infty} \frac{\text{Tr}_{Adj} P^n}{n^4} + \frac{M^2 T^2}{2\pi^2} \sum_{n=1}^{\infty} \frac{\text{Tr}_{Adj} P^n}{n^2}. \quad (64)$$

It is important to note that the mass parameter M should not be interpreted as a gauge boson mass, nor do we limit ourselves to $ML \ll 1$: the additional term in V_g is purely phenomenological. The crucial feature of this potential is that for sufficiently large values of the dimensionless parameter M/T , the second term dominates and the potential leads to a $Z(N)$ -symmetric, confining minimum for P . On the other hand, for small values of M/T , the first term dominates and the pure gauge theory will be in the deconfined phase. This model has the nice property that V_G is a good representation of the gauge boson contribution for high temperatures.

Minimizing V_G as a function of the eigenvalues of P leads to expressions for the pressure and associated thermodynamic quantities. In the case of $SU(3)$, we can set the mass scale M by requiring that V_g yields the correct deconfinement temperature for the pure gauge theory, with a value of $T_d = 270 \text{ MeV}$, giving $M = 596 \text{ MeV}$. The pressure p is given by the value of $-V_G$ at the minimum of the potential. The behavior obtained for the pressure, the energy density ϵ , and the interaction measure $\Delta \equiv (\epsilon - 3p)/T^4$ are all roughly consistent




Figure 4: The Polyakov loop $Tr_F P$ as a function of T in $SU(3)$ using a phenomenological model [42, 71].

with lattice simulations for $T > T_d$ and can be improved with the use of one or two additional free parameters [69, 70]. As shown in Fig. 4, the order parameter TrP for the deconfinement transition jumps at T_d , indicating a first-order deconfinement transition for $SU(3)$.

C. Role of quarks and other particles

1. Chiral symmetry breaking

If confinement is the most striking feature of QCD, chiral symmetry breaking is certainly a close second. If we imagine for a moment a variant of QCD in which all N_f quarks have the same mass, there will be a global $SU(N_F)$ symmetry which rotates the different flavors into each other. In this theory, the left- and right-handed components of the Dirac fields, defined by

$$q_{L/R} = \frac{1}{2}(1 \pm \gamma_5)q \quad (65)$$

are connected by the Lagrangian mass term

$$m\bar{q}q = m(\bar{q}_L q_R + \bar{q}_R q_L) \quad (66)$$

which can be thought of as an interaction that flips the helicity of a quark. If the masses of the quarks are set to zero, then the left- and right-handed components are no longer coupled, and the symmetry group becomes $SU(N_f)_L \otimes SU(N_f)_R$. At the classical level, there is also a $U(1)_A$ axial symmetry generated by γ_5 , but it is anomalous.

In our universe, the two lightest quarks, the u and d quarks, have masses m_u and m_d which are more than an order of magnitude smaller than Λ_{QCD} , so we are close to having an $SU(2)_L \otimes SU(2)_R$ symmetry. On the basis of models and, more recently, lattice simulations, we believe that in the massless limit this symmetry would be spontaneously broken to $SU(2)$, the symmetry group of the massive theory, by the existence of a condensate $\langle 0 | \bar{u}u + \bar{d}d | 0 \rangle$. The breaking of this continuous global symmetry leads to the existence of three Goldstone bosons, the pions.

Unfortunately, our ability to understand chiral symmetry breaking analytically has been limited. We assume that the basic mechanism is similar to the mechanism of superconductivity. In the case of superconductivity, electron-hole interactions mediated by phonons lead to an expectation value for a composite field, while in gauge theories, the gauge boson interactions somehow give rise to some effective four-fermion interaction which leads to the formation of a $\bar{q}q$ condensate. However, there is no convincing derivation of this effective interaction from the gauge theory. In the absence of a more fundamental understanding, Nambu-Jona Lasinio (NJL) models are often used to describe the chiral symmetry effects. In NJL models, a four-fermion interaction induces chiral symmetry breaking; the four-fermion interaction is non-renormalizable, and NJL models cannot be regarded as fundamental. There has been a great deal of work on NJL models, both as phenomenological models for hadrons and as effective theories of QCD [72, 73].

NJL models use purely fermionic interactions as a proxy for the gauge theory interactions that give rise to chiral symmetry breaking. Typically, the relation between the gauge theory and an associated NJL model is fixed by matching important hadronic parameters such as f_π . In the case of NJL models at finite temperature, and particularly PNJL models, it is common to assume that the NJL model parameters are fixed by $T = 0$ hadronic parameters and remain constant as T is increased, at least up to the deconfinement temperature.

As an example of an NJL model consider the representative model [71]

$$L_{NJL} = \bar{\psi} (i\gamma \cdot \partial - m_0) \psi + \frac{g_S}{2} \left[(\bar{\psi} \lambda^a \psi)^2 + (\bar{\psi} i\gamma_5 \lambda^a \psi)^2 \right] + g_D [\det \bar{\psi} (1 - \gamma_5) \psi + h.c.] \quad (67)$$

where ψ is associated with N_f flavors of Dirac fermions in the fundamental or adjoint representation of the gauge group $SU(N)$. The λ^a 's are the generators of the flavor symmetry group $U(N_f)$; g_S represents the strength of the four-fermion scalar-pseudoscalar coupling and g_D fixes the strength of an anomaly induced term. For simplicity, we can take the mass matrix m_0 to be diagonal: $(m_0)_{jk} = m_{0j}\delta_{jk}$. If m_0 and g_D are taken to be zero, L_{NJL} is invariant under the global symmetry $U(N_f)_L \times U(N_f)_R$. With a nonzero, flavor-independent mass m_0 and $g_D \neq 0$, the symmetry is reduced to $SU(N_f)_V \times U(1)_V$. This model can be solved approximately in a self-consistent manner. The effective potential V_{F0} can be calculated at zero temperature as a function of the chiral condensates $\sigma_j \equiv \bar{\psi}_j \psi_j$:

$$V_{F0}(m_j) = \sum_j g_S \sigma_j^2 + 2g_D (N_f - 1) \prod_j \sigma_j - 2N \sum_{j=1}^{N_f} \int \frac{d^3k}{(2\pi)^3} \omega_k^{(j)} \quad (68)$$

where the last term representing the sum of fermionic zero-point energies from the functional determinant. The zero-point energy of each mode $\omega_k^{(j)} = \sqrt{k^2 + m_j^2}$, is written in terms of the constituent mass m_j , given in this model as

$$m_j \equiv m_j^0 - 2g_S \sigma_j - 2g_D (N_f - 1) \prod_{k \neq j} \sigma_k. \quad (69)$$

The sum over zero-point energies is divergent, and requires regularization. Differentiation with respect to σ_j yields a set of gap equations which can be solved numerically for the chiral condensate.

2. Quarks and deconfinement

The addition of quarks or other particles to a gauge theory can completely change the theory's finite temperature behavior. Such particles can directly affect deconfinement, because they can alter or even destroy center symmetry. These effects can be seen directly in the effective potential.

The calculation of the one-loop effective potential for gauge bosons can be extended to other particles such as quarks in a very general way [74]. Because the effective potential is central for subsequent discussion, we treat the calculation in detail. All of the required expressions can be obtained from the one-loop effective potential for a charged scalar boson of mass M moving in a background $U(1)$ Polyakov loop in d spatial dimensions. This is

given by

$$V_B(\theta) = \frac{1}{\beta} \int \frac{d^d k}{(2\pi)^d} \ln [1 - e^{-\beta\omega_k + i\theta}] + \frac{1}{\beta} \int \frac{d^d k}{(2\pi)^d} \ln [1 - e^{-\beta\omega_k - i\theta}] \quad (70)$$

where ω_k is given by $\sqrt{k^2 + M^2}$. Now consider a scalar boson in the fundamental representation of an $SU(N)$ gauge group moving in a uniform Polyakov loop background. The one-loop contribution to the finite-temperature effective potential is

$$V = \frac{1}{\beta} \int \frac{d^d k}{(2\pi)^d} \text{Tr}_F [\ln (1 - P e^{-\beta\omega_k}) + \ln (1 - P^\dagger e^{-\beta\omega_k})] \quad (71)$$

where the first term is due to particles and the second term is due to antiparticles. A global unitary transformation puts P into the diagonal form

$$P_{jk} = \delta_{jk} \exp(i\theta_j) \quad (72)$$

and the partition function can be written as

$$V = \sum_j V_B(\theta_j). \quad (73)$$

As a second example, consider the case of the gauge bosons themselves, which lie in the adjoint representation of the gauge group. The Polyakov loop in the adjoint representation is an $(N^2 - 1) \times (N^2 - 1)$ matrix. The partition function for the $N^2 - 1$ particles is

$$s \frac{1}{2} \sum_{j,k=1}^N (1 - \frac{1}{N} \delta_{jk}) V_B(\theta_j - \theta_k) \quad (74)$$

where the δ_{jk} removes a singlet contribution, and the factor of $1/2$ corrects for over-counting since V_B has both a particle and antiparticle contribution. The factor s accounts for spin degeneracy; in $3+1$ dimensions $s = 2$, a consequence of the two possible polarization states of gauge bosons.

For our third and final example, consider the evaluation of fermionic partition functions, which can be reduced to the general bosonic problem. A typical fermionic contribution of particle and antiparticle has the form

$$V_F(\theta) = -\frac{1}{\beta} \int \frac{d^d k}{(2\pi)^d} \ln [1 + e^{-\beta\omega_k + i\theta}] - \frac{1}{\beta} \int \frac{d^d k}{(2\pi)^d} \ln [1 + e^{-\beta\omega_k - i\theta}] \quad (75)$$

which is easily written as

$$V_F(\theta) = -V_B(\pi + \theta). \quad (76)$$

For fermions in the fundamental representation of $SU(N)$, the partition function is

$$s \sum_j V_F(\theta_j) = -s \sum_j V_B(\pi + \theta_j) \quad (77)$$

where the factor s again accounts for spin degeneracy.

A low-temperature expansion for $V_B(\theta)$ can be generated for arbitrary spatial dimension d by expanding the logarithm and integrating term by term:

$$V_B(\theta) = -\frac{M^{d/2+1/2}}{2^{d/2-3/2}\pi^{d/2+1/2}\beta^{d/2+1/2}} \sum_{n=1}^{\infty} \frac{1}{n^{d/2+1/2}} K_{(d+1)/2}(n\beta M) \cos(n\theta) \quad (78)$$

which gives for scalar bosons in the fundamental representation

$$V_B(\theta) = -\frac{M^{d/2+1/2}}{2^{d/2-1/2}\pi^{d/2+1/2}\beta^{d/2+1/2}} \sum_{n=1}^{\infty} \frac{1}{n^{d/2+1/2}} K_{(d+1)/2}(n\beta M) \text{Tr}_F(P^n + P^{\dagger n}) \quad (79)$$

with similar results for bosons in other representations. For fermions, we have

$$V_F(\theta) = \frac{M^{d/2+1/2}}{2^{d/2-3/2}\pi^{d/2+1/2}\beta^{d/2+1/2}} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^{d/2+1/2}} K_{(d+1)/2}(n\beta M) \cos(n\theta). \quad (80)$$

In a path integral representation, the factors of $(-1)^n$ are a consequence of fermionic antiperiodic boundary conditions.

A high-temperature expansion for $V_B(\theta)$ may also be obtained in $3+1$ dimensions

$$V_B(\theta) = -\frac{2}{\pi^2\beta^4} \left[\frac{\pi^4}{90} - \frac{1}{48}\theta_+^4 + \frac{\pi}{12}\theta_+^3 - \frac{\pi^2}{12}\theta_+^2 \right] + \frac{M^2}{2\pi^2\beta^2} \left[\frac{1}{4}\theta_+^2 - \frac{\pi}{2}\theta_+ + \frac{\pi^2}{6} \right] \quad (81)$$

$$- \frac{1}{2\pi\beta^4} \sum_{l \in \mathbb{Z}} R(\beta M, \theta, l) - \frac{M^4}{16\pi^2} \left[\ln \left(\frac{\beta M}{4\pi} \right) + \gamma - \frac{3}{4} \right]$$

where θ_+ is θ modulo 2π such that $0 \leq \theta < 2\pi$. The sum $\sum_l$ over l is over all integers with

$$R(\beta M, \theta, l) \equiv \frac{1}{3} [(\beta M)^2 + (\theta - 2\pi l)^2]^{3/2} - \frac{1}{3} |\theta - 2\pi l|^3 - \frac{1}{2} |\theta - 2\pi l| \beta^2 M^2 - \frac{(\beta M)^4}{16\pi |l|} \quad (82)$$

except that the divergent term is omitted when $l = 0$.

The first term in eqn. 81 is the blackbody free energy for two degrees of freedom, and depends only on the temperature and the angle θ . The second term, which is the leading correction due to the mass M at high temperatures, often appears in discussions of symmetry restoration at high temperatures with $\theta = 0$. The third term is closely associated with the $n = 0$ Matsubara mode, which is the most infrared singular contribution to a finite temperature functional determinant. This term is responsible for non-analytic behavior in

finite temperature perturbation theory via the summation of ring diagrams. For example, in a scalar theory it gives rise to the $\lambda^{3/2}$ contribution to the free energy; in QED, the contribution is e^3 [75]. The last term is logarithmic in the dimensionless combination βM and independent of θ . In calculations of effective potentials, it typically combines with zero-temperature logarithms in such a way that the temperature T sets the scale of running coupling constants at high T .

3. Lattice results

Physical QCD has neither center symmetry, because there are quarks, not chiral symmetry, because there is a quark mass term in the Lagrangian. Nevertheless, simulations of $SU(3)$ gauge theories with quarks have revealed a complicated phase structure, where the number of flavors and the quark masses are crucial parameters. Quark masses span several orders of magnitude, from the light u and d quarks, with $u, d \ll \Lambda_{QCD}$ to the very heavy t quark with $m_t \gg \Lambda_{QCD}$. Because m_s is roughly on the order of Λ_{QCD} , lattice simulations aimed at realistic descriptions of hadronic physics are generally carried out with $2+1$ flavors for sea quarks, that is, two light flavors and one intermediate relative to Λ_{QCD} .

For gauge theories at finite temperature, the static approximation is the next logical step beyond pure gauge theories, and represents very well the effect of very heavy particles. In the case where quarks or other particles in the fundamental representation satisfy $\beta M \gg 1$, we need keep only the leading term in the low-temperature expansion

$$V_H = -h_F [Tr_F P + Tr_F P^\dagger]. \quad (83)$$

This generalizes immediately to particles in a representation R . Recalling the spin model interpretation, we see that a heavy particle behaves like an external field h_R coupled to the Polyakov loop $Tr_R P$ in the representation R . For fermions in the fundamental representation, this breaks $Z(N)$ symmetry explicitly. In the case of $SU(3)$ and other theories with first-order deconfinement transitions, the addition of a small $Z(N)$ symmetry-breaking field does not remove the transition, as it would in the case of a second-order transition. Instead, a small symmetry-breaking field with $h_F > 0$ makes the deconfined phase more favorable at the deconfinement temperature, and there is a new deconfinement temperature, giving rise to a critical line in the $T - h_F$ plane. On both sides of the line, $Tr_F P$ is non-zero, but

there is a discontinuity that grows smaller with increasing h_F . This behavior persists until a critical value of h_F is reached, where the first-order line terminates in a critical end point, believed to be in the usual ϕ^4 universality class [76–78].

As quark masses are lowered, chiral symmetry effects become important. Phase transitions, if present, lie at temperatures below the deconfinement transition temperature of the pure gauge theory, and the phase structure is very sensitive to quark masses. The results of the extensive lattice studies that have been done is usually displayed in a so-called Columbia plot, which shows the phase structure of three-flavor QCD, with up, down and strange quarks, by showing lines of phase transitions as functions of a common up and down quark mass m_{ud} and the strange quark mass m_s . A version of this plot is shown in Figure 5. There are several interesting limiting cases contained in this diagram:

- $m_{ud} = m_s = \infty$: Because the quarks are completely removed from the dynamics, this limit is a pure gauge theory. For $SU(3)$, the phase transition is first-order and associated with the spontaneous breaking of center symmetry, as described above. If the quark masses are decreased from infinity, center symmetry is explicitly broken, but the phase transition persists for large values of the quark masses, manifesting as a jump in $\langle Tr_F P \rangle$ from one non-zero value to another. As explained above, this region of first-order transitions terminates in what appears in the figure as a second order line.
- $m_{ud} = 0 ; m_s = \infty$ This is the two-flavor chiral limit of QCD, with a second-order transition in the $\mathcal{O}(4)$ universality class that extends down from $m_s = \infty$ until it meets another critical line at a tricritical point.
- $m_{ud} = m_s = 0$ This is the three-flavor chiral limit of QCD. Of course, it is less realistic than the two-flavor limit because the strange quark is sufficiently heavy to be on the order Λ . This is a first-order transition that extends into the $m_{ud} - m_s$ plane before terminating in a second-order line.

Physical QCD does not appear to have a finite temperature phase transition. Nevertheless, remnants of both the deconfinement transition of the pure gauge theory and the chiral transition of massless quarks remain. Near the critical temperature of the chiral transition in the massless theory, there is a crossover region. This is a relatively narrow range of




Figure 5: Schematic version of the Columbia plot showing phase structure as a function of the m_{ud} and m_s quark masses

temperatures over which p , ϵ , $\bar{\psi}\psi$ and $Tr_F P$ change rapidly. In physical QCD, this crossover marks the change from low-temperature hadronic behavior to high-temperature quark-gluon plasma behavior.

4. connection of chiral symmetry breaking and confinement

If a system has two or more order parameters, their behavior is typically correlated. This can be understood as arising from terms in the effective potential allowed by symmetry that couple the order parameters. In general, a first-order transition observed as a discontinuity in one order parameter will also give rise to a discontinuity in others. In the case of QCD and related theories, lattice simulations with fundamental representation fermions show that deconfinement and chiral symmetry restoration are strongly correlated.

The connection of chiral symmetry breaking and deconfinement can be seen in strong-coupling lattice models [79]. Integration over the spatial link variables does two things: generates the nearest-neighbor Polyakov loop coupling that drives the deconfinement transition in strong coupling, and produces a four-fermion coupling that leads to spontaneous chiral symmetry breaking at low T . In order to explore the interrelationship of confinement and chiral symmetry breaking in a continuum model, we can use a generalization of Nambu-Jona Lasinio models known as Polyakov-Nambu-Jona Lasinio (PNJL) models [80]; see [81] for an alternative approach. NJL models have been used to study hadronic physics at finite temperature, but they include only chiral symmetry restoration, and do not model deconfinement. This omission is rectified by the PNJL models, which include both chiral restoration and deconfinement. In PNJL models, fermions with NJL couplings move in a nontrivial Polyakov loop background, and the effects of gluons at finite temperature is modeled in a semi-phenomenological way.

The key idea of PNJL models is familiar: all particles move in a constant Polyakov loop background, and the effective potential will be a function of both the chiral order parameter $\bar{\psi}\psi$ and P . The coupling between the chiral condensate and the Polyakov loop arises because the covariant derivative D_μ replaces the conventional derivative ∂_μ in the fermion kinetic term. However, D_μ contains only a constant A_4 gauge field background that gives rise to a non-trivial Polyakov loop. The complete PNJL effective potential consists of three terms

$$V_{PNJL} = V_G(P) + V_{F0}(\sigma_j) + V_{FT}(P, \sigma_j) \quad (84)$$

where $V_G(P)$ reproduces the behavior of the pure gauge theory, $V_{F0}(\sigma_j)$ is the NJL effective potential at $T = 0$, and $V_{FT}(P, \sigma_j)$ is the $T \neq 0$ part of the fermion effective potential, given by

$$V_{FL}(P, m) = -2 \sum_j Tr_F [T \int \frac{d^3k}{(2\pi)^3} \ln(1 + P e^{-\beta \omega_k^{(j)}}) + h.c.] \quad (85)$$

where the $\omega_k^{(j)}$'s are defined in terms of the constituent masses as they were for the NJL model in Section III C. If V_G is given by eqn. 64 and V_{F0} by 68, then minimizing V_{PNJL} as a function of σ for two flavors with $m_0 = 5.5 \text{ MeV}$ gives the behavior shown in Fig. 6 [71]. This shows the explanatory power of PNJL models. The constituent mass m is heavy at low temperatures, due to chiral symmetry breaking. The larger the constituent mass, the smaller the $Z(3)$ breaking effect of the fermions, reflected in the small value of $\langle Tr_F P \rangle$ at




Figure 6: Order parameters in a PNJL model of QCD with two light flavors as a function of temperature [71].

low temperatures. On the other hand, a small value for $\langle Tr_F P \rangle$ reduces the effectiveness of finite-temperature effects in restoring chiral symmetry. These synergistic effects combine in the case of fundamental representation fermions to give a single crossover temperature at which both order parameters are changing rapidly, consistent with the rapid crossover seen in lattice simulations.

Chiral and deconfinement behavior are not always as closely correlated as they are with quarks in the fundamental representation. In the case of adjoint fermions, lattice simulations have shown that chiral symmetry is restored at a substantially higher temperature than the deconfinement temperature, with the ratio of critical temperatures T_c/T_d around 7.8 [82, 83]. This behavior can also be reproduced in adjoint PNJL models [71].

D. finite density QCD

The phase diagram of QCD in the temperature-chemical potential plane is of great interest. Unfortunately, lattice gauge theory is not directly applicable because of the sign problem. The nature of the problem can be seen from the one-loop effective potential. The effective potential $V_{FT\mu}$ for a single flavor quarks at temperature $T = \beta^{-1}$ and chemical

potential μ in a static Polyakov loop background P is

$$V_{FT\mu} = -2T \int \frac{d^3k}{(2\pi)^3} Tr_F [\ln (1 + P e^{\beta\mu - \beta\omega_k}) + \ln (1 + P^\dagger e^{-\beta\mu - \beta\omega_k})] \quad (86)$$

reflecting the fact that a non-zero chemical potential behaves as an imaginary $U(1)$ Polyakov loop. This expression is complex for $\mu \neq 0$ when P is non-trivial. This can be seen very clearly for heavy quarks, where $V_{FT\mu}$ can be written as

$$V_{FT\mu} \approx f_q \approx -h_F [e^{\beta\mu} Tr_F P + e^{-\beta\mu} Tr_F P^\dagger]. \quad (87)$$

Because $Tr_F P$ is complex for $N \geq 3$, the effective action for the gauge fields is complex. This is one version of the sign problem for gauge theories at finite density: the Euclidean path integral involve complex weights. This problem is a fundamental barrier to lattice simulations of QCD at finite density [84, 85] and a significant problem in other fields [86].

A large part of the interest in finite-density QCD lies in the possibility of a color superconducting phase of hadronic matter in the interior of neutron stars [16]. At non-zero density, there are additional order parameters reflecting the possibility of Cooper pair formation whenever an attractive channel exists between fermions. As we have seen, chiral symmetry breaking can be viewed as the formation of a condensate of $\bar{q}q$ pairs in the vacuum. Other bifermion condensates are possible when the chemical potential is non-zero, giving rise to what is known as color superconductivity. This is an extension of the BCS mechanism from the formation of Cooper pairs of electrons to qq condensates. The mechanism for BCS superconductivity is an attractive force between Cooper pairs mediated by phonons. Early work on color superconductivity considered both interactions mediated by gluon exchange and induced by instantons via the axial $U(1)$ anomaly. In practice, it is common to assume an NJL model with a set of four-fermion interactions, perhaps with an anomaly-induced term as well that is a $2N_f$ fermion coupling.

Because quarks carry both color and flavor, it is necessary to consider both when discussing appropriate order parameters for color superconductivity. Quarks carry flavor, color and spinor indices, so the quark condensate in a given channel can be characterized by

$$\langle \psi_{ia}^\alpha \psi_{jb}^\beta \rangle = P_{ijab}^{\alpha\beta} \Delta \quad (88)$$

where α and β are color indices, j and k are flavor indices, and a and b are spinor indices. The color-flavor-spin matrix characterizes a particular pairing, and the gap parameter Δ gives the

magnitude of the condensate in this channel. The condensate may be momentum-dependent. The familiar BCS condensate is a special case where the condensate is independent of position, so the pairing is between fermions of equal and opposite momentum, and a spin singlet. Unlike the fermion-antifermion condensates, the fermion-fermion condensates are not gauge invariant, and thus are similar to the Higgs scalar expectation value, which similarly depends on the choice of gauge. As with the Higgs mechanism, physical observables like the spectrum are gauge-invariant. There are strong arguments why the color-flavor locking phase (CFL) phase is favored for high-density QCD, combining antisymmetry in color and flavor with a Lorentz scalar behavior, giving

$$\left\langle \psi_i^\alpha C \gamma_5 \psi_j^\beta \right\rangle \propto \epsilon^{\alpha\beta A} \epsilon_{ijA} \quad (89)$$

where C is the charge-conjugation matrix acting on Dirac spinors. For a recent review of our current understanding of the phase diagram of dense QCD, see [87].

IV. PHASES OF GAUGE THEORIES ON $R^3 \times S^1$

In the last few years, it has proven possible to construct four-dimensional gauge theories for which confinement may be reliably demonstrated using semiclassical methods [1, 2]. These models combine $Z(N)$ symmetry, the effective potential for P , instantons, and monopoles into a satisfying picture of confinement for a special class of models. All of the models in this class have one or more small compact directions. Models with an $R^3 \times S^1$ topology have been most investigated, and discussion here will focus on this class. The use of one or more compact directions will cause the running coupling constant of an asymptotically free gauge theory to be small, so that semiclassical methods are reliable. For example, if the circumference L of S^1 is small, *i.e.*, $L \ll \Lambda^{-1}$, then $g(L) \ll 1$. However, this leads to an immediate problem: generally speaking, one or more small compact directions lead to breaking of $Z(N)$ symmetry in those directions. As we have seen in Section III B, for the case of finite temperature gauge theories, where $L = \beta = 1/T$, the effective potential for the Polyakov loop is easily calculated to lowest order in perturbation theory; for a pure gauge theory it is given by eq. (56):

$$V_{gauge}(P, \beta) = \frac{-2}{\pi^2 \beta^4} \sum_{n=1}^{\infty} \frac{Tr_{Adj} P^n}{n^4} \quad (90)$$

where the trace of P in the adjoint representation is given by $Tr_A P = |Tr_F P|^2 - 1$. This effective potential is minimized when $P = zI$ where $z \in Z(N)$, indicating that $Z(N)$ symmetry is spontaneously broken at high temperatures where the one-loop expression is valid. However, it is possible to maintain $Z(N)$ symmetry even when L is small by modifying the action. This leads to a perturbative calculation of possible phase structures, which turns out to be very rich, as well as a perturbative understanding of Polyakov loop physics in the confined phase. Furthermore, the restoration of $Z(N)$ symmetry for small L leads to a non-perturbative mechanism for confinement, as measured by Wilson loops orthogonal to the compact direction. In this confinement mechanism, a key role is played by finite-temperature instantons, also known as calorons, and their monopole constituents. Thus we obtain a realization of a long-held scenario for quark confinement, based on ideas originally proposed by Mandelstam [88, 89] and 't Hooft [90, 91].

A. Restoring center symmetry via deformations or periodic adjoint fermions

There are two broad approaches to maintaining $Z(N)$ symmetry for small L . The first approach deforms the pure gauge theory by adding additional terms to the gauge action [2, 92, 93]. The general form for such a deformation is

$$S \rightarrow S + \beta \int d^3x \sum_{k=1}^{\infty} a_k Tr_A P(\vec{x}, x_4)^k \quad (91)$$

where the value of x_4 is arbitrary and can be taken to be 0. Such terms are often referred to as double-trace deformations; see Fig. 7. If the coefficients a_k are sufficiently large, they will counteract the effects of the one-loop effective potential, and $Z(N)$ symmetry will hold for small L . Strictly speaking, only the first $[N/2]$ terms are necessary to ensure confinement. As discussed in Section IIIB, it is easy to prove that for a classical Polyakov loop P , the conditions $Tr_F P^k = 0$ with $1 \leq k \leq [N/2]$ determine the unique set of Polyakov loop eigenvalues that constitute a confining solution, *i.e.*, one for which $Tr_R P = 0$ for all representations with $k_R \neq 0$ [42]. The explicit solution is simple: up to a factor necessary to ensure $\det P = 1$, the eigenvalues of P are given by the set of N 'th roots of unity, which are permuted by a global $Z(N)$ symmetry transformation. The effective potential associated




Figure 7: A double-trace Polyakov loop on $R^3 \times S^1$.

with S is given approximately by

$$V_{eff}(P, \beta) = \frac{-2}{\pi^2 \beta^4} \sum_{n=1}^{\infty} \frac{Tr_{Adj} P^n}{n^4} + \sum_{k=1}^{\left[\frac{N}{2}\right]} a_k Tr_{Adj} P^k. \quad (92)$$

For sufficiently large and positive values of the a_k 's, the confined phase yields the lowest value of V_{eff} . However, a rich phase structure emerges from the minimization of V_{eff} for intermediate values of the coefficients a_k . For $N \geq 3$, the effective potential predicts that one or more phases may separate the deconfined phase from the confined phase. In the case of $SU(3)$, a single new phase is predicted, and has been observed in lattice simulations [2]. For larger values of N , there is a rich set of possible phases, including some where $Z(N)$ breaks down to a proper subgroup $Z(p)$. In such phases, particles in the fundamental representation are confined, but bound states of p particles are not [92].

Lattice simulations of $SU(3)$ and $SU(4)$ agree for small L with the theoretical predictions based on effective potential arguments [2]. The phase diagram of $SU(3)$ as a function of $T = L^{-1}$ and a_1 has three phases: the confined phase, the deconfined phase, and a new phase, the skewed phase. In general, the three phases of the eigenvalues of the Polyakov loop may be taken to be the set $\{\theta_1, \theta_2, \theta_3\}$ where $\theta_1 + \theta_2 + \theta_3 = 0$. For all three phases, it is possible to use $Z(3)$ symmetry to make $Tr_F P$ real, and reduce the phases to the set $\{0, \theta, -\theta\}$ such that $Tr_F P = 1 + 2 \cos \theta$. The deconfined phase is represented by $\theta = 0$,

the confined phase is given by $\theta = 2\pi/3$, and the skewed phase by $\theta = \pi$. An important result obtained from the lattice simulation of $SU(3)$ is that the small- L confining region, where semiclassical methods yield confinement, are smoothly connected to the conventional large- L confining region. In the case of $SU(4)$, a sufficiently large value of a_1 leads to a partially-confining phase where $Z(4)$ is spontaneously broken to $Z(2)$. Particles with $k = 1$ are confined in this phase, *i.e.*, $\langle Tr_F P(\vec{x}) \rangle = 0$, but particles with $k = 2$ are not, as indicated by $\langle Tr_F P^2(\vec{x}) \rangle \neq 0$. As in the case of $SU(3)$, $Tr_F P$ can be made real. Perturbation theory then predicts a deconfined phase where the phases of the eigenvalues of P are $\{0, 0, 0, 0\}$, a confined phase where they are $\{\pi/4, 3\pi/4, 5\pi/4, 7\pi/4\}$, and a partially confined phase where the phases are $\{\pi/2, \pi/2, -\pi/2, -\pi/2\}$. The confined phase corresponds to the matrix P_1 and the partially confined phase to the matrix P_2 described in Section III B.

Another approach to preserving $Z(N)$ symmetry for small L uses fermions in the adjoint representation with periodic boundary conditions in the compact direction [1]. In this case, it would be somewhat misleading to use β as a synonym for L , because the transfer matrix for evolution in the compact direction is not positive-definite. Periodic boundary conditions in the compact direction imply that the generating function of the ensemble, *i.e.*, the partition function, is given by

$$Z = Tr \left[(-1)^F e^{-LH} \right] \quad (93)$$

where F is the fermion number and H is the Hamiltonian in the compact direction. This graded ensemble, familiar from supersymmetry, can be obtained from an ensemble $Tr [\exp(\beta\mu F - \beta H)]$ with chemical potential μ by the replacement $\beta\mu \rightarrow i\pi$. This system can be viewed as a gauge theory with periodic boundary conditions in one compact spatial direction of length $L = \beta$, and the transfer matrix in the time direction is positive-definite,

The use of periodic boundary conditions for the adjoint fermions dramatically changes their contribution to the Polyakov loop effective potential. In perturbation theory, the replacement $\beta\mu \rightarrow i\pi$ shifts the Matsubara frequencies from $\beta\omega_n = (2n + 1)\pi$ to $\beta\omega_n = 2n\pi$. The one loop effective potential is now essentially that of a bosonic field, but with an overall negative sign due to fermi statistics [74]. The sum of the effective potential for the fermions plus that of the gauge bosons gives

$$V_{eff}(P, \beta, m, N_f) = \frac{1}{\pi^2 \beta^4} \sum_{n=1}^{\infty} \frac{Tr_{Adj} P^n}{n^2} \left[2N_f \beta^2 m^2 K_2(n\beta m) - \frac{2}{n^2} \right] \quad (94)$$

where N_f is the number of adjoint Dirac fermions and m is their mass [49]. Note that the

first term in brackets, due to the fermions, is positive for every value of n , while the second term, due to the gauge bosons, is negative.

The largest contribution to the effective potential at high temperatures is typically from the $n = 1$ term, which can be written simply as

$$\frac{1}{\pi^2 \beta^4} [2N_f \beta^2 m^2 K_2(\beta m) - 2] [|Tr_F P|^2 - 1] \quad (95)$$

where the overall sign depends only on N_f and βm . If $N_f \geq 1$ and βm is sufficiently small, this term will favor $Tr_F P = 0$. On the other hand, if βm is sufficiently large, a value of P from the center, $Z(N)$, is preferred. Note that an $\mathcal{N} = 1$ super Yang-Mills theory would correspond to $N_f = 1/2$ and $m = 0$, giving a vanishing perturbative contribution for all n [94, 95]. In that case, non-perturbative effects lead to a confining effective potential for all values of β . In the case of $N_f \geq 1$, each term in the effective potential will change sign in succession as m is lowered towards zero. For larger values of N , this leads to a cascade of phases separating the confined and deconfined phases [48]. Numerical investigation shows that the confined phase is obtained if $N\beta m \lesssim 4.00398$ [49]. As m increases, it becomes favorable that $Tr_F P^n \neq 0$ for successive values of n . If N is even, the first phase after the confined phase will be a phase with $Z(N/2)$ symmetry. As m increases, the last phase before reaching the deconfined phase will have $Z(2)$ symmetry, in which $k = 1$ states are confined, but all states with higher k are not. Lattice simulations of $SU(3)$ with periodic adjoint fermions are completely consistent with the picture [96] predicted by the effective potential, with a skewed phase separating the confined phase and deconfined phase. For $N \geq 3$, there are generally phases intermediate between the confined and deconfined phases which are not of the partially-confined type. Careful numerical analysis appears to be necessary on a case-by-case basis to determine the phase structure for each value of N [48]. There has been very interesting work on lattice gauge theories in the large- N limit that indicates the same rich phase structure [97, 98]; this work will be discussed in Section VI.

There are some interesting additional issues arising when periodic adjoint fermions are used to obtain $Z(N)$ symmetry for small L . There are strong indications from strong-coupling lattice calculations [79] and the closely related Polyakov-Nambu-Jona Lasinio (PNJL) models [80] that the mass m that appears in the effective potential $V(P)$ should be regarded as a constituent mass that includes the substantial effects of chiral symmetry breaking. In strong-coupling lattice calculations and PNJL models, this effect is responsible

for the coupling of P and $\bar{\psi}\psi$ in simulations of QCD at finite temperature. Thus it is important that simulations of $SU(3)$ with $N_f = 2$ flavors of adjoint fermions show explicitly that the $Z(N)$ -invariant confined phase is regained when the fermion mass is sufficiently small [96]. However, this raises another issue. For a simple double-trace deformation, the small- L and large- L regions are smoothly connected. A semi-phenomenological analysis based on a PNJL model [71] suggests that this is also the case with periodic adjoint fermions, but additional modifications of the action may be necessary to realize the connection.

B. Monopoles and instantons on $R^3 \times S^1$

The non-perturbative dynamics of confining gauge theories on $R^3 \times S^1$ are based on Polyakov's analysis of the Georgi-Glashow model in three dimensions [99]. This is an $SU(2)$ gauge model coupled to an adjoint Higgs scalar. The models we are considering thus differ by the addition of a fourth compact dimension and a change to the action designed to maintain $Z(N)$ symmetry. The four-dimensional Georgi-Glashow model is the standard example of a gauge theory with classical monopole solutions when the Higgs expectation value is non-zero. These monopoles make a non-perturbative contribution to the partition function Z . In three dimensions, these monopoles are instantons. Polyakov showed that a gas of such three-dimensional monopoles gives rise to non-perturbative confinement in three dimensions, even though the theory appears to be in a Higgs phase perturbatively.

Because L is small in the $R^3 \times S^1$ models we consider, the three-dimensional effective theory describing the behavior of Wilson loops in the non-compact directions will have many features in common with the three-dimensional theory first discussed by Polyakov. In the four-dimensional theory, monopole solutions with short worldline trajectories in the compact direction exist, and behave as three-dimensional instantons in the effective theory; see Fig. 8. In models on $R^3 \times S^1$, the role of the three-dimensional scalar field is played by the fourth component of the gauge field A_4 . In a gauge where the Polyakov loop is diagonal and independent of x_4 , P has a vacuum expected value induced by the perturbative effective potential. However, there is another way to understand the presence of monopoles in this phase, based on studies of instantons in pure gauge theories at finite temperature and the properties of the KvBLL caloron solution [100–102]. If the Polyakov loop has a non-trivial expectation value, finite-temperature instantons in $SU(N)$ may be decomposed




Figure 8: Short monopole worldline on $R^3 \times S^1$.

into N monopoles, and the locations of the monopoles become parameters of the moduli space of the instanton. In the case of $SU(2)$, an instanton may be decomposed into a conventional BPS monopole and a so-called KK (Kaluza-Klein) monopole. The presence of the KK monopole solution differentiates the case of a gauge field at finite temperature from the case of an adjoint scalar breaking $SU(N)$ to $U(1)^{N-1}$, in which case there are $N - 1$ fundamental monopoles. We will consider in detail the simplest case of $N = 2$.

The BPS monopole is found using the standard arguments [103, 104]. The Euclidean Lagrangian $\mathcal{L}$ can be taken to be

$$\mathcal{L} = \frac{1}{4} (F_{\mu\nu})^2 + V_{eff}(P) \quad (96)$$

where V_{eff} includes both the one-loop gluonic effective potential and the additional term that prevents $Z(N)$ symmetry breaking. This can also be written as

$$\mathcal{L} = \frac{1}{2} (D_j A_4)^2 + \frac{1}{2} (B_j)^2 + V_{eff}(P). \quad (97)$$

We can associate with $\mathcal{L}$ an energy defined by

$$E = \int d^3x \left[\frac{1}{2} (B_j)^2 + \frac{1}{2} (D_j A_4)^2 + V_{eff}(P) \right] \quad (98)$$

as well as an action $S = LE$. We will concern ourselves for now with the solutions in the BPS limit, in which the effective potential V_{eff} is neglected, but the boundary condition on

P at infinity imposed by the potential is retained. We can write the energy as

$$E = \int d^3x \left[\frac{1}{2} (B_j \pm D_j A_4)^2 \mp B_j D_j A_4 \right]. \quad (99)$$

This expression is a sum of squares plus a term which can be converted to a surface integral, giving rise to the BPS inequality

$$E \geq \mp \int dS_j B_j A_4. \quad (100)$$

The BPS inequality is saturated if the equality $B_j = \mp D_j A_4$ holds. For the case of a single monopole at the origin, we require the fields at spatial infinity to behave as

$$\begin{aligned} \lim_{r \rightarrow \infty} A_4^a &= w \frac{x^a}{r} \\ \lim_{r \rightarrow \infty} A_i^a &= \epsilon^{aij} \frac{x_j}{gr^2}. \end{aligned} \quad (101)$$

Note that w is related to the eigenvalues of P at large distances by $w = 2\theta/gL$. Note that A_4 has the usual hedgehog form. A_i^a is chosen such that covariant terms vanish at infinity: $(D_i A_4)^a = 0$. With the 't Hooft-Polyakov ansatz, the general expressions for the fields become

$$\begin{aligned} A_4^a &= wh(r) \frac{x^a}{r} \\ A_i^a &= a(r) \epsilon^{aij} \frac{x_j}{gr^2} \end{aligned} \quad (102)$$

where we define $w > 0$ and require $h(\infty) = 1$ or -1 , and $a(\infty) = 1$ to obtain the correct asymptotic behavior. We must also have $h = a = 0$ at $r = 0$ to have well-defined functions at the origin. We identify a magnetic flux

$$\Phi = \pm \int dS_j B_j^a \frac{x^a}{r} = \mp \frac{4\pi}{g} \quad (103)$$

where the $+$ sign corresponds to the case $h(\infty) = 1$ and $-$ corresponds to $h(\infty) = -1$. The energy of the BPS monopole can be written as

$$E_{BPS} = \mp \Phi w = \frac{4\pi w}{g}. \quad (104)$$

In addition to the BPS monopole, there is another, topologically distinct monopole which occurs at finite temperature when A_4 is treated as a Higgs field [94]. Starting from a static monopole solution where $|A_4| = w$ at spatial infinity, we apply a special gauge transformation

$$U_{special} = \exp \left[-\frac{i\pi x_4}{L} r^3 \right] \quad (105)$$

where τ^i is the Pauli matrix. $U_{special}$ transforms A_μ in such a way that the value of A_4 at spatial infinity is shifted: $w \rightarrow w - 2\pi/gL$. If we instead start from a static monopole solution such that $A_4 = 2\pi/gL - w$ at spatial infinity, then the action of $U_{special}$ gives a monopole solution with $A_4 = -w$ at spatial infinity. A final constant gauge transformation $U_{const} = \exp[i\pi\tau^2/2]$ yields a new monopole solution with $A_4 = w$ at spatial infinity. The distinction between the BPS solution, which is independent of x_4 , and the KK solution is made clear by consideration of the topological charge. The action of $U_{special}$ followed by U_{const} increases the topological charge by 1 and changes the sign of the monopole charge. Thus the KK solution is topologically distinct from the BPS solution because it carries instanton number 1. The $\overline{BPS}$ antimonopole has magnetic charge opposite to the BPS monopole, and hence the same as that of the KK monopole. The $\overline{KK}$ monopole has the same magnetic charge as the BPS monopole, but carries instanton number -1 . This is all completely consistent with the KvBLL decomposition of instantons in the pure gauge theory with non-trivial Polyakov loop behavior, where $SU(2)$ instantons can be decomposed into a BPS monopole and a KK monopole. Our picture of the confined phase is one where instantons and anti-instantons have “melted” into their constituent monopoles and anti-monopoles, which effectively forms a three-dimensional gas of magnetic monopoles. In the BPS limit, both the magnetic and scalar interactions are long-ranged; this behavior appears prominently, for example, in the construction of N -monopole solutions in the BPS limit.

The BPS solution has action

$$S_{BPS} = \frac{4\pi wL}{g} = \frac{8\pi\theta}{g^2}. \quad (106)$$

For the KK solution, we have instead

$$S_{KK} = \frac{4\pi(2\pi - gLw)}{g^2} = \frac{4\pi(2\pi - 2\theta)}{g^2}. \quad (107)$$

The sum $S_{BPS} + S_{KK}$ is exactly $8\pi^2/g^2$, the action of an instanton. For $\theta = \pi/2$, the $Z(2)$ -symmetric value for $SU(2)$, $S_{BPS} = S_{KK}$. This extends to $SU(N)$, where the action of a monopole of any type is $8\pi^2/g^2N$.

Although we used the BPS construction to exhibit the existence and some properties of the monopole solutions of our system, we must move away from the BPS limit to ensure that magnetic interaction dominate at large distances, *i.e.*, that the three-dimensional scalar interactions associated with A_4 and are not long-ranged. This behavior is natural in the

confined, where the characteristic scale of the Debye (electric) screening mass associated with A_4 is large, on the order of g/L . It is well known that the BPS bound for the monopole mass holds as an equality only when the scalar potential is taken to zero. Numerical studies [105] have shown that the monopole action is given in general for $SU(2)$ as

$$LE_{BPS}C(\epsilon) \tag{108}$$

where C a function of the quartic term in the potential that varies from $C = 1$ in the BPS limit to a maximum value $C(\infty) = 1.787$. Thus corrections to the BPS result for the monopole mass and action due to the potential terms are less than a factor of two. We will henceforth use the exact results for the actions in the BPS limit, neglecting corrections from V_{eff} for the sake of simplicity of notation.

The $SU(2)$ construction of BPS and KK monopoles extends to $SU(N)$ in the standard way, via the embedding of $SU(2)$ subgroups in $SU(N)$. There are $N - 1$ BPS monopoles and 1 KK monopole inside an instanton. In the confined phase, each of the N monopoles has action $8\pi^2/g^2N$. It has long been thought that instanton effects must be suppressed in the large- N limit, because instanton effects would vanish as $\exp(-cN)$ in the limit $N \rightarrow \infty$ with $\lambda \equiv g^2N$ fixed [106]. In contrast, we see that the effects of monopole constituents of instantons are not suppressed by the large- N limit.

C. Monopoles and space-like string tensions

It is important to understand that Wilson loops in planes orthogonal to the compact direction should show area law behavior, even if the $Z(N)$ symmetry associated with the compact direction is broken. This is an old observation about the deconfined phase [107, 108] which is very clearly observed in lattice simulations of $SU(2)$ and $SU(3)$ at temperatures above the deconfinement transition [109, 110]. At first sight, this seems to directly conflict with the association of deconfinement with the loss of area-law behavior for Wilson loops. However, the introduction of a compact direction, as in the case of finite temperature, explicitly breaks space-time symmetry. In the case of finite temperature, Wilson loops measuring electric flux have perimeter behavior in the deconfined phase; Wilson loops measuring magnetic flux still obey an area law. This asymmetry in behavior can be understood on the basis of center symmetry. The full center symmetry of an $SU(N)$ gauge theory on a d -dimensional




Figure 9: Spatial Wilson loop in $R^3 \times S^1$ geometry.

hypertorus T^d is $Z(N)^d$. While the $Z(N)$ symmetry may break spontaneously in the short compact direction, the other $Z(N)$ symmetries are unbroken, and thus the associated Wilson loops obey an area law.

In order to understand the effects of monopoles play in the confined phase, we must analyze their interactions. We begin with a discussion of quantum fluctuations around the monopole solutions. The contribution to the partition function of a single BPS monopole at finite temperature was considered by Zarembo [111]. The measure factor $d\mu^a$ associated with the collective coordinates (moduli) of the monopole solution, including the Jacobians from the zero modes is given by [95]

$$\int d\mu^a = \mu^4 \int \frac{d^3x}{(2\pi)^{3/2}} J_x \int_0^{2\pi} \frac{d\phi}{(2\pi)^{1/2}} J_\phi \quad (109)$$

where x is the position and ϕ the $U(1)$ phase of the monopole and μ is a Pauli-Villars regulator. The label a denotes the type of monopole, $a = \{BPS, KK, \overline{BPS}, \overline{KK}\}$. The Jacobians are

$$J_x = S_a^{3/2}, \quad J_\phi = NL S_a^{1/2}. \quad (110)$$

Each of the four zero modes contributes a factor of μ . In the BPS limit, each monopole carries an overall factor

$$\begin{aligned} Z_a &= c\mu^{7/2} (NL)^{1/2} S_a^2 \exp[-S_a + \mathcal{O}(1)] \int d^3x \\ &= \xi_a \exp[-S_a] \int d^3x \end{aligned} \quad (111)$$

in its contribution to Z [111]. The factor ξ_a is $c\mu^{7/2}(NL)^{1/2}S_a^2$ where c is a numerical constant and the factor of d^3x represents the integration over the location of the monopole. From the construction of the KK monopole, we see that we have $\xi_{KK}(\theta) = \xi_{BPS}(\pi - \theta)$.

The renormalization of the functional determinant arising from quantum fluctuations around the monopole solution is particularly simple in the confined phase, as first observed by Davies *et al.* in the corresponding supersymmetric model [94]. The dependence on the Pauli-Villars regulator is removed, as usual, by coupling constant renormalization. The relation at one loop of the bare coupling and the regulator mass μ to a renormalization-group invariant scale Λ is

$$\Lambda^{b_0} = \mu^{b_0} e^{-8\pi^2/g^2 N} \quad (112)$$

where b_0 is the first coefficient of the β function divided by N :

$$b_0 = \frac{11}{3} - \frac{4}{3} \cdot \frac{n_f C(R_f)}{N} - \frac{1}{6} \cdot \frac{n_b C(R_b)}{N} \quad (113)$$

where n_f is the number of flavors of Dirac fermions in a representation R_f , n_b is the number of flavors of real scalars in a representation R_b , and $C(R)$ is obtained from $Tr_R(T^a T^b) = C(R)\delta^{ab}$. For the case of a pure gauge theory with a deformation, there are four collective coordinates and this gives a factor of μ^4 . The functional integral over gauge degrees of freedom gives rise to a factor $\det'[-D^2]^{-1} \propto \mu^{-1/3}$ and the action contributes a factor $\exp(-8\pi^2/g^2 N)$ in the confined phase. Thus the contribution of a single monopole to the partition function gives a factor

$$\mu^{4-\frac{1}{3}} e^{-8\pi^2/g^2 N} = \mu^{11/3} e^{-8\pi^2/g^2 N} = \Lambda^{11/3}. \quad (114)$$

Thus detailed calculation confirms what we might have guessed on dimensional grounds: the contribution $\xi_a e^{-8\pi^2/g^2 N} \propto L^{-3} (\Lambda L)^{11/3}$. Note that the eliminations of renormalization-dependent quantities by renormalization-independent quantities depends crucially on the coefficient of $1/g^2$ in the action.

The interaction of the monopoles is essentially the one described by Polyakov in his original treatment of the Georgi-Glashow model in three dimensions [99], slightly generalized to include both the BPS and KK monopoles. Let us consider, say, a BPS-type monopole and KK-type monopole located at $\vec{x}_1$ and $\vec{x}_2$ in the non-compact directions, with static worldlines in the compact direction. The interaction energy due to magnetic charge of such

a pair is

$$E_{BPS-KK} = - \left(\frac{4\pi}{g} \right)^2 \frac{1}{4\pi |\vec{x}_1 - \vec{x}_2|} \quad (115)$$

and the associated action is approximately $S_{BPS} + S_{KK} + LE_{BPS-KK}$. As discussed above, this will be larger than the value obtained from the Bogomolny bound, but of the same order of magnitude. There is an elegant way to capture the dynamics of the monopole plasma, using an Abelian scalar field σ dual to the magnetic field. Assuming that the Abelian magnetic gauge field is three-dimensional for small L , we may write

$$L \int d^3x \frac{1}{2} B_k^2 = \int d^3x \frac{g^2}{32\pi^2 L} (\partial_k \sigma)^2 \quad (116)$$

where the normalization of σ is chosen to simplify the form of the interaction terms. The three-dimensional effective action is given by

$$L_{eff} = \frac{g^2}{32\pi^2 L} (\partial_j \sigma)^2 - \sum_a \xi_a e^{-S_a + i q_a \sigma} \quad (117)$$

where the sum is over the set $\{BPS, KK, \overline{BPS}, \overline{KK}\}$. Each species of monopole has its own magnetic charge sign $q_a = \pm$ as well as its own action S_a . The coefficients ξ_a represent the functional determinant associated with each kind of monopole, but the combination $\xi_a \exp(-S_a)$ may be usefully regarded as a monopole activity in terms of the statistical mechanics of a gas of magnetic charges. The generating functional

$$Z_\sigma = \int [d\sigma] \exp \left[- \int d^3x L_{eff} \right] \quad (118)$$

is precisely equivalent to the generating function of the monopole gas. This equivalence may be proved by expanding Z_σ in a power series in the ξ_a 's, and doing the functional integral over σ for each term of the expansion.

The magnetic monopole plasma leads to confinement in three dimensions. For our effective three-dimensional theory, any Wilson loop in a hyperplane of fixed x_4 , for example a Wilson loop in the $x_1 - x_2$ plane, will show an area law; see figure 9. The original procedure of Polyakov [99] may be used to calculate the string tension, where the presence of a large planar Wilson loop causes the dual field σ to have a discontinuity on the surface associated with the loop and a half-kink profile on both sides. However, an alternative procedure is simpler where the discontinuity in the gauge field strength induced by the Wilson loop is moved to infinity so that the string tension is obtained from the kink solution connecting the two vacua of the dual field σ [93].

In the confined phase, the action and functional determinant factors for all four types of monopoles are the same, so we denote them by S_M and ξ_M . The potential term in the mixed and confined phases then reduces to

$$-\sum_a \xi_a e^{-S_a + i q_a \sigma} \rightarrow 4 \xi_M e^{-S_M} [1 - \cos(\sigma)] \quad (119)$$

which has minima at $\sigma = 0$ and $\sigma = 2\pi$; we have added a constant for convenience such that the potential is positive everywhere and zero at the minima. A one-dimensional soliton solution $\sigma_s(z)$ connects the two vacua, and the string tension σ_{3d} for Wilson loops in the three non-compact directions is given by

$$\sigma_{3d} = \int_{-\infty}^{+\infty} dz L_{eff}(\sigma_z(z)) \quad (120)$$

which can be calculated via a Bogomolny inequality to be

$$\sigma_{3d} = \frac{4g}{\pi} \sqrt{\frac{\xi_M}{L}} e^{-S_M}. \quad (121)$$

Of course, this result depends on L and cannot be used outside the region $\Lambda L \ll 1$; nevertheless, this is a concrete realization of confinement in a four-dimensional field theory via non-Abelian monopoles.

D. The special case of supersymmetry

The earliest indication that confinement on $R^3 \times S^1$ was interesting and tractable came about in an unlikely way, in an effort to resolve a controversy over adjoint fermion condensates in four-dimensional $\mathcal{N} = 1$ supersymmetric $SU(N)$ gauge theories. Davies *et al.* resolved the controversy by considering supersymmetric $SU(N)$ gauge theories on $R^3 \times S^1$ [94, 95]. In such theories, supersymmetry requires that the number of adjoint fermion degrees of freedom (“gluinos”) equals the number of gauge degrees of freedom, and that the adjoint fermions have the same boundary conditions as the gauge fields. This model interpolates between four-dimensional $\mathcal{N} = 1$ supersymmetric $SU(N)$ gauge theories as $L \rightarrow \infty$ and three-dimensional $\mathcal{N} = 2$ supersymmetric $SU(N)$ gauge theories in the limit $L \rightarrow 0$ with no intervening phase transition as L is varied. In the three-dimensional limit, the scalar field of the $d = 3$ $\mathcal{N} = 2$ theory is simply the $n = 0$ Matsubara mode of A_4^3 . We follow closely the original notation and treatment of $SU(2)$ by Davies *et al.* [94], but see also [1, 112]; some

subtleties of the calculation that were previously unappreciated are discussed in detail in [113]. As before, a non-vanishing expectation value for A_4^3 breaks the $SU(2)$ symmetry to $U(1)$, and the low-energy effective theory will have only $U(1)$ symmetry. The superpotential of this model is trivial in perturbation theory and the effective potential is completely non-perturbative. The superpotential can be written in terms of chiral and anti-chiral $\mathcal{N} = 1$ superfields

$$\begin{aligned}\Phi &= Z + \sqrt{2}\theta^\alpha\Psi_\alpha + \theta^\alpha\theta_\alpha\mathcal{F} \\ \bar{\Phi} &= \bar{Z} + \sqrt{2}\bar{\theta}^\alpha\bar{\Psi}_\alpha + \bar{\theta}^\alpha\bar{\theta}_\alpha\bar{\mathcal{F}}\end{aligned}$$

where Z is a complex scalar $\phi + i\gamma$ that includes both the three-dimensional scalar $\phi = A_4^3$ and a field γ which is dual to the $U(1)$ magnetic field B_j^3 and thus proportional to σ in our notation. The effective action is

$$S_{eff} = S_{cl} + \frac{L}{g^2} \int d^3x \left[\int d^2\theta \mathcal{W}(\Phi) + \int d^2\bar{\theta} \bar{\mathcal{W}}(\bar{\Phi}) \right] \quad (122)$$

where $\mathcal{W}(\Phi)$ is given by

$$\mathcal{W}(\Phi) = \left(\frac{g^2}{4\pi L} \right)^2 M \left[\exp\left(-\frac{4\pi L}{g^2}\Phi\right) + \exp\left(-\frac{8\pi^2}{g^2} + \frac{4\pi L}{g^2}\Phi\right) \right]. \quad (123)$$

The parameter M is $M = 16\pi^2 L^2 M_{PV}^3$ with M_{PV} the Pauli-Villars regulator associated with the functional determinant. The effective potential is given by

$$V_{eff} = \bar{\mathcal{F}}\mathcal{F} = \frac{\partial \mathcal{W}}{\partial Z} \frac{\partial \bar{\mathcal{W}}}{\partial \bar{Z}} \quad (124)$$

with

$$\mathcal{F} = \frac{\partial \mathcal{W}}{\partial Z} = -\frac{Mg^2}{4\pi L} \left[\exp\left(-\frac{4\pi L}{g^2}\Phi\right) - \exp\left(-\frac{8\pi^2}{g^2} + \frac{4\pi L}{g^2}\Phi\right) \right] \quad (125)$$

The minimum of V_{eff} preserves supersymmetry: $\mathcal{F}(\langle Z \rangle = 0)$ implies $\langle \phi \rangle = \pi/\beta$ and $\langle \gamma \rangle = 0$, and leads to $Tr_F P = 0$. The effective potential for γ is then given by

$$V_{eff}(\phi = \pi/\beta, \gamma) = 2 \left(\frac{Mg^2}{4\pi L} \right)^2 \exp\left(-\frac{8\pi^2}{g^2}\right) \left[1 - \cos\left(\frac{8\pi L}{g^2}\gamma\right) \right] \quad (126)$$

The string tension is again obtained from the one-dimensional kink solution of the equation of motion obtained from the effective action for the dual field. This model represents a lower boundary of where periodic adjoint fermions work to restore $Z(N)$ symmetry, with

the superpartner of the gauge boson corresponding to $N_f = 1/2$ for the number of Dirac fermions. For $N_f = 1/2$, the one-loop effective potential Eqn. 94 favors the deconfined phase for when the fermion mass is positive, and vanishes identically when it is zero.

E. Polyakov loop string tensions

Polyakov loop string tensions are calculable perturbatively in the small- L confining region from small fluctuations about the confining minimum of the effective potential [49, 114]; see also [93]. The effective Lagrangian can be written in terms of the phases of the Polyakov loop eigenvalues θ_j

$$\frac{1}{g^2} \sum_{j=1}^N (\nabla \theta_j)^2 + V_{eff}(\theta) \quad (127)$$

where the kinetic term is obtained from the standard kinetic term for gauge theories and the potential term is the one-loop effective potential. In the high- T (small- L) confining region, the minimum of the effective potential will have $Z(N)$ symmetry. A convenient form for such a solution is

$$(P_0)_{jk} = w z^j \delta_{jk} \quad (128)$$

where $z = \exp(2\pi i/N)$ and w is a phase factor that ensures $\det P_0 = 1$. For small fluctuations, we write

$$P = P_0 e^{i\delta\theta} \quad (129)$$

where $\delta\theta$ lies in the Cartan algebra of $SU(N)$, *i.e.*, is a diagonal traceless matrix, and therefore has $N - 1$ independent components. The moments of the Polyakov loop are

$$Tr_F P^k = w^k \sum_{j=1}^N z^{jk} e^{ik\delta\theta_j} \quad (130)$$

If k is not divisible by N , we have approximately

$$Tr_F P^k \simeq i k w^k \sum_{j=1}^N z^{jk} \delta\theta_j = i k w^k \delta\tilde{\theta}_k \quad (131)$$

where $\delta\tilde{\theta}$ is the discrete Fourier transform of $\delta\theta$, related by

$$\delta\tilde{\theta}_k = \sum_{j=1}^N z^{jk} \delta\theta_j \quad (132)$$

$$\delta\theta_j = \frac{1}{N} \sum_{k=1}^N z^{-jk} \delta\tilde{\theta}_k. \quad (133)$$

Note that the reality of $\delta\theta$ implies that $\delta\tilde{\theta}_k^* = \delta\tilde{\theta}_{N-k}$; the last Fourier component, $\delta\tilde{\theta}_N$, is identically zero, due the tracelessness of $\delta\phi$. If k is divisible by N we have instead

$$Tr_F P^k \simeq N - \frac{1}{2} k^2 \sum_{j=1}^N (\delta\theta_j)^2 = N - \frac{1}{2N} k^2 \sum_{m=1}^N \left(\delta\tilde{\theta}_m \delta\tilde{\theta}_{N-m} \right). \quad (134)$$

These formulae allow us to write the three-dimensional effective action in terms of the $\delta\tilde{\theta}_k$ to quadratic order. For each value of k , the terms in the potential which contribute have $n \equiv k \bmod N$, $n \equiv N - k \bmod N$, or $n \equiv 0 \bmod N$. We obtain a different mass σ_k^P/T for each Fourier component $\delta\tilde{\theta}_k$ measured by using Polyakov loops of N -ality k . There are $[N/2]$ different string tensions because $\sigma_k^P = \sigma_{N-k}^P$. The string tensions obtained, for example, in lattice simulations will depend on the operators used. For example, the operators $(Tr_F P)^k \sim \left(\delta\tilde{\theta}_1 \right)^k$ and $Tr_F(P^k) \sim \delta\tilde{\theta}_k$ both have N -ality k , but the two-point correlation function of the first operator will decay as $\exp[-k\sigma_1^P r/T]$, while the second operator decays as $\exp[-\sigma_k^P r/T]$. Group characters will in general have a complicated pattern of mixing. Consider the symmetric/anti-symmetric representations of dimension $N(N \pm 1)/2$, with characters given by

$$\chi_{S/A}(P) = (1/2) \left[(Tr_F P)^2 \pm Tr_F P^2 \right]. \quad (135)$$

These operators will have two different contributions, corresponding to $2\sigma_1^{(t)}$ and $\sigma_2^{(t)}$. At higher orders in perturbation theory, there will be mixing, and ultimately only the lightest state for a given N -ality will be seen at large distances. This phenomena, often referred to as string breaking, was first observed in lattice simulations in $SU(2)$ Higgs models [115–117] and later seen in pure $SU(2)$ gauge theory in $(2+1)$ dimensions [118].

The behavior seen in the models considered in this section is very different from the behavior seen in exactly solvable two-dimensional gauge theories or in lattice simulations of four-dimensional gauge theories. This is not surprising. For the case of double-trace deformations, there are at least $[N/2]$ independent parameters which can be adjusted, and which can vary in turn the σ_k^P . For the case of periodic adjoint fermion, the results are

continuous functions of βm . The string tensions are of order g :

$$\begin{aligned} \left(\frac{\sigma_k^P}{T}\right)^2 &= g^2 N \frac{N_f m^2}{\pi^2} \sum_{j=0}^{\infty} [K_2((k+jN)\beta m) + K_2((N-k+jN)\beta m) - 2K_2((j+1)N\beta m)] \\ &\quad - g^2 N \frac{T^2}{3N^2} \left[3 \csc^2\left(\frac{\pi k}{N}\right) - 1 \right] \end{aligned} \quad (136)$$

where the gluon contribution has been summed in the last term. Note that the symmetry $\sigma_k^P = \sigma_{N-k}^P$ is manifest in this formula. The $m = 0$ limit has the simple form

$$\left(\frac{\sigma_k^P}{T}\right)^2 = \frac{(2N_f - 1) g^2 T^2}{3N} \left[3 \csc^2\left(\frac{\pi k}{N}\right) - 1 \right] \quad (137)$$

and is a good approximation for $\beta m \ll 1$. This scaling law is not at all like either Casimir or sine-law scaling, because the usual hierarchy $\sigma_{k+1}^P \geq \sigma_k^P$ is here reversed. Because we expect on the basis of $SU(3)$ simulations that the high-temperature confining region is continuously connected to the conventional low-temperature region, there must be an inversion of the string tension hierarchy between the two regions for all $N \geq 4$. Other features of the string tension behavior are in line with our expectations. For the case $N_f = 1/2$, corresponding to a single multiplet of adjoint Majorana fermions, the perturbative string tension vanishes, and it is the non-perturbative contribution to the effective potential induced by monopoles that gives rise to the string tension in this case [94, 95]. The large- N limit of Eqn. 137 is smooth. For fixed k as $N \rightarrow \infty$, we have

$$\left(\frac{\sigma_k^P}{T}\right)^2 \sim \frac{(2N_f - 1) \lambda T^2}{\pi^2 k^2} \quad (138)$$

where λ is the 't Hooft coupling $g^2 N$. The Polyakov loop string tensions obtained in the small- L confining region, while conforming to general principles, do not appear to tell us much about confinement at large L .

F. Other geometries

It is possible to apply the same methods used for gauge theories on $R^3 \times S^1$ to other geometries, such as $S^1 \times S^3$ or $R^2 \times T^2$. Meyers and Hollowood have performed a detailed study of $SU(N)$ gauge theories on $S^1 \times S^3$ with periodic adjoint fermions [119]. In this geometry, R^3 is replaced by S^3 , so there are two length scales introduced by the geometry, the radius of the three-sphere $R = R_{S^3}$ and $L = R_{S^1}$. We require $\min[R_{S^1}, R_{S^3}] \ll \Lambda$ so that

we are in the weak-coupling region. The projection onto gauge-invariant states, manifested as integration over the eigenvalues of the Polyakov loop, ensures non-trivial behavior. Because the spatial volume is finite, there is no actual phase transition for finite N , only a crossover as R/L is varied. However, the large- N limit does give a phase transition whose behavior is closely approximated even for moderate values of N .

As in the case of $R^3 \times S^1$, the information about the phase structure is contained in the effective potential. Each field adds a factor like

$$\pm \text{Tr}_R \log [-D_0^2 - \Delta] \quad (139)$$

to the effective potential, where Δ is the appropriate Laplacian on S^3 for each kind of field, $\pm$ for bosonic and fermionic fields. With N_f flavors of fermion, the effective potential is given by

$$S(P) = \sum_{n=1}^{\infty} \frac{1}{n} \left\{ (1 - z_v(nL/R)) \text{Tr}_{\text{Adj}}(P^n) + \sum_{f=1}^{N_f} z_f(nL/R, m_f R) \text{Tr}_F(P^n) \right\} \quad (140)$$

where the first term is due to Haar measure. The second term is from the gauge boson, and as expected takes the form of a sum over paths winding around S^1 , with z_v given by

$$z_v(L/R) = 2 \sum_{\ell=1}^{\infty} \ell(\ell+2) e^{-L(\ell+1)/R} = \frac{6e^{-2L/R} - 2e^{-3L/R}}{(1 - e^{-L/R})^3}. \quad (141)$$

The final term, due to the fermions, has a similar interpretation, with

$$z_f(L/R, mR) = 2 \sum_{\ell=1}^{\infty} \ell(\ell+1) e^{-L\sqrt{(\ell+1/2)^2 + m^2 R^2}/R}. \quad (142)$$

Although the fermionic contribution cannot be put into a simple closed form, Hollowood and Myers derive the useful form

$$z_f\left(\frac{L}{R}, mR\right) = \frac{2m^2 R^3}{L} K_2(Lm) - \frac{mR}{2} K_1(Lm) + 4 \int_{mR}^{\infty} dx \frac{x^2 + \frac{1}{4}}{e^{2\pi x} + 1} \sin(L\sqrt{x^2 - m^2 R^2}/R) \quad (143)$$

In the limit $m \rightarrow 0$, this reduces to

$$z_f(L/R, 0) = \frac{4e^{-3L/2R}}{(1 - e^{-L/R})^3} \equiv \sum_{\ell=1}^{\infty} 2\ell(\ell+1) e^{-L(\ell+1/2)/R} \quad (144)$$

and in the limit $R \rightarrow \infty$ with fixed m and L , we recover

$$z_f(L/R, mR) \longrightarrow \frac{2m^2 R^3}{L} K_2(Lm) \quad (145)$$

which is the expression that one obtains by working directly on $R^3 \times S^1$ [48].

In the case of adjoint fermions, we have

$$S(P) = \sum_{n=1}^{\infty} \frac{1}{n} \left(1 - z_v(nL/R) + N_f z_f(nL/R, m_f R) \right) \sum_{ij=1}^N \cos(n(\theta_i - \theta_j)) . \quad (146)$$

In the large- N limit, the effective potential becomes a functional of the distribution $\rho(\theta)$ of Polyakov loop eigenvalues, with

$$S[\rho(\theta)] = N^2 \int d\theta \int d\theta' \rho(\theta) \rho(\theta') \sum_{n=1}^{\infty} \frac{f(nL/R, mR)}{n} \cos(n(\theta - \theta')) \quad (147)$$

where

$$f(L/R, mR) = 1 - z_B(L/R) + N_f z_F(L/R, mR) \quad (148)$$

and ρ is normalized to

$$\int_0^{2\pi} d\theta \rho(\theta) = 1 . \quad (149)$$

This expression for $S[\rho(\theta)]$ can be made even simpler by writing ρ in terms of its Fourier components

$$\rho(\theta) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} \rho_n e^{in\theta} \quad (150)$$

where $\rho_n = \text{Tr}_F P^n / N$. Then we may write the effective potential as

$$S[\rho(\theta)] = \frac{N^2}{2} \sum_{n=1}^{\infty} \frac{f(nL/R, mR)}{n} |\rho_n|^2 \quad (151)$$

It is now obvious that the effective potential will become unstable in the contribution of N -ality n precisely when $f(nL/R, mR) = 0$. These techniques can also be applied to the study of gauge theories at finite temperature and density on $S^3 \times S^1$ [120–122]. Another interesting geometry is $R^2 \times T^2$, where $SU(N)$ gauge theories are in the universality class of $Z(N) \times Z(N)$ spin models because there are two compact directions [123, 124]. Dimensional reduction leads to two-dimensional models where many powerful techniques are available, as discussed in Section V.

G. Higgs theories on $R^3 \times S^1$

As shown by 't Hooft [90, 91], there is a fundamental conflict between the Higgs mechanism and confinement. The dual superconductor picture of confinement gives a simple

picture of this conflict as being between an electric condensate in the Higgs mechanism and a magnetic condensate in a confining model. With analytic control of confinement in gauge theory on $R^3 \times S^1$, an adjoint Higgs field can be introduced to study the interplay of confinement and the Higgs mechanism analytically [125, 126]. The simplest case is $SU(2)$, with a classical Euclidean action given by

$$S_c = \int d^4x \left[\frac{1}{4} (F_{\mu\nu}^a)^2 + \frac{1}{2} (D_\mu \phi)^T \cdot D_\mu \phi + V(\phi) \right] \quad (152)$$

where the Higgs potential is

$$V(\phi) = \frac{1}{2} m^2 \phi^2 + \frac{1}{4} \lambda (\phi^2)^2 \quad (153)$$

with $\phi^2 = \phi^T \phi$. The action has a $Z(2)_H$ global symmetry given by $\phi \rightarrow -\phi$, in addition to $Z(2)_C$ center symmetry which transform P to $-P$.

The scalar field ϕ is not gauge invariant, and cannot serve as an order parameter for the breaking of the $Z(2)_H$ symmetry associated with ϕ when gauge interactions are present. This is an old problem, a consequence of Elitzur's theorem [127]. Higgs models with scalar fields in the fundamental and adjoint representations behave differently. For Higgs models with scalar fields in the fundamental representation, the confined and Higgs phases are connected [128], in a manner similar to the connection between liquid and gas phases. In this case, center symmetry is explicitly broken, and large Wilson loops do not have area-law behavior due to screening by the scalars. In the adjoint case, center symmetry is preserved by the action, and there is a distinct phase transition between the confined and Higgs phases. On $R^3 \times S^1$, there are three distinct gauge-invariant order parameters associated with the $Z(2)_C \times Z(2)_H$ symmetry. The first of these is the trace in the fundamental representation of the Polyakov loop P itself, $\langle \text{Tr}_F P \rangle$. It transforms non-trivially under $Z(2)_C$ but is invariant under $Z(2)_H$. The second is $\langle \text{Tr}_F [P^2(x) \phi(x)] \rangle$ which is invariant under $Z(2)_C$, but transforms non-trivially under $Z(2)_H$. Finally, there is $\langle \text{Tr}_F [P(x) \phi(x)] \rangle$, which transforms non-trivially under both groups.

Using the three order parameters, one can show that there are four distinct phases: a deconfined phase, a confined phase, a Higgs phase, and a mixed confined phase. The mixed confined phase occurs where one might expect a phase in which there is both confinement and the Higgs mechanism, but the behavior of the order parameters distinguishes the two phases. In the mixed confined phase, the $Z(2)_C \times Z(2)_H$ global symmetry breaks spontaneously to a

$Z(2)$ subgroup that acts non-trivially on both the scalar field and the Polyakov loop. In the mixed confined phase, the role of the scalar field is played a linear combination of the Higgs field ϕ and A_4 in the construction of BPS and KK monopole solutions. In all four phases, Wilson loops orthogonal to the compact direction are expected to show area-law behavior due to unbroken center symmetry in the non-compact directions. This confining behavior can be attributed to a dilute monopole gas in a broad region that includes portions of all four phases.

The supersymmetric analog of this model is the Seiberg-Witten model [129], which is an $\mathcal{N} = 2$ supersymmetric gauge theory with gauge group $SU(2)$. Seiberg and Witten found that in this model the addition of an $\mathcal{N} = 1$ mass perturbation leads to confinement by magnetic monopoles. Recently, Poppitz and Unsal have examined the behavior of this model on $R^3 \times S^1$, and concluded that the confined phase seen for small compactification circumference on $R^3 \times S^1$ is connected to the confining phase at infinite compactification circumference [130]. In their work, Euclidean monopoles in which a linear combination of A_4 and ϕ plays the role of the scalar field appear in a very similar fashion to the non-supersymmetric model.

V. CONFORMALITY AND DUALITY

A. Non-trivial fixed points in $d = 3 + 1$ gauge theories

Up until this point, we have implicitly restricted ourselves to theories like QCD which have a simple renormalization group structure: they have an ultraviolet fixed point at $g = 0$, and hence are asymptotically free, and an infrared fixed point at $g = \infty$ when the theory is defined on R^4 . Gauge theories with non-trivial fixed point structure are known to occur if the number of particle representations included in the theory is sufficiently large. The first two coefficients of the perturbative contribution to the renormalization-group β function of a gauge theory are independent of the renormalization scheme, and given by

$$\beta(g^2) = \frac{dg^2}{d \log q^2} = -\frac{b_1}{16\pi^2}g^4 - \frac{b_2}{(16\pi^2)^2}g^6 + \dots \quad (154)$$

where

$$b_1 = \frac{11}{3} C_2(G) - \frac{4}{3} N_f T(R)$$

$$b_2 = \frac{34}{3} [C_2(G)]^2 - N_f T(R) \left[\frac{20}{3} C_2(G) + 4C_2(R) \right]$$

The coefficients b_1 and b_2 depend on the group and the group representations of the particles via $C_2(R)$, the value of the quadratic Casimir operator in representation R (G denotes the adjoint representation, so $C_2(G) = N_c$), while $T(R)$ is the normalization of the group generators in R : $Tr(T_R^a T_R^b) = T(R) \delta^{ab}$.

Three possible behaviors may be obtained from perturbation theory:

- 1) $b_1 < 0$: the theory is no longer asymptotically free at high energies, and $g = 0$ is an infrared fixed point, as is the case in QED. For a non-Abelian gauge theory, this is typically the case when the matter content of the theory becomes too large. Only the gauge fields themselves give a negative contribution to b_1 , so a sufficient number of additional matter fields can drive b_1 negative.
- 2) $b_1 > 0$ and $b_2 > 0$: in this case the coupling runs out of the region of perturbative computability in the IR. This is the QCD-like case, where the theory is asymptotically free at high energies (the UV) and confining at low energies (the IR).
- 3) $b_1 > 0$ and $b_2 < 0$: Perturbation theory predicts a non-trivial fixed point, given by

$$g_{BZ}^2 = -16\pi^2 \frac{b_1}{b_2}. \quad (155)$$

These three behaviors are shown in 10. The corresponding phase structure as a function of the number of flavors is shown in figure 11. As was first pointed out by Banks and Zaks [131], this fixed point may or may not be located in the region where perturbation theory is reliable. However, when it is located in that region, the prediction of an infrared-stable fixed point is reliable. Note however that a change in the renormalization prescription, which redefines the coupling constant, will change the beta function. This in turn will change the location of a non-trivial fixed point. Given that such theories are possible, we would like to understand their properties. Such theories are not generally candidates for describing the real world. Theories with non-trivial infrared fixed points have correlation functions




Figure 10: Three possible behaviors for the beta function. If $N_f < N_f^*$, the theory is asymptotically free but has no non-trivial fixed point. If $N_f > N_f^{AF}$, the gauge theory is no longer asymptotically free. For $N_f^* < N_f < N_f^{AF}$, there is a conformal window with a non-trivial infrared fixed point.

that decay algebraically at large distances, in a manner familiar from critical behavior at second-order phase transitions. Requiring that models not have an infrared fixed point is a condition that likely requires non-perturbative understanding that lattice gauge theories can supply. The case of a non-trivial infrared fixed point, case 3), is intermediate between the familiar case 2) and non-asymptotically free theories. Typically, for a given gauge group and choice of additional representation R , there is some number of flavors N_f^{AF} such that asymptotic freedom is lost, and some smaller value N_f^* where an infrared fixed point appears. Theories with $N_f^* < N_f < N_f^{AF}$ are said to lie in the conformal window.

There is a close relation between the existence of the conformal window and possible beyond the standard model physics associate with the Higgs field. In the standard model, the Higgs field is a fundamental scalar field, in the sense of not being a composite; in fact it is the only fundamental scalar field in the standard model. Because the bare Higgs mass receives corrections which are quadratic in any ultraviolet cutoff, there is a problem of “naturalness” or “fine-tuning” in achieving a sufficiently small Higgs mass. This problem




Figure 11: Phase structure of a theory with a conformal window as a function of the number of flavors N_f .

would be circumvented if the Higgs were a bound state of some new fermions rather than a fundamental scalar. Theories of this type that have a non-fundamental Higgs are called technicolor models, and the gauge group of the new fermions is the technicolor group; see [132] for a recent review. In the standard model, the Higgs is responsible not just for the gauge boson masses, but also for the fermion masses. The class of models needed for successful phenomenology are called extended technicolor models, and seem to require a gauge theory very close to the conformal window. Such theories are often referred to as “walking” because the running coupling constant moves slowly if the beta function is small, as it would be near a zero.

There has been much work done on the location of the conformal window but definitive results have not yet been achieved. Much of the analytic work uses the chiral order parameter $\bar{\psi}\psi$ as the order parameter for the transition from a confining phase, where $\bar{\psi}\psi$ is expected to be non-zero, to a conformal phase which has no dynamically generated scale. One method is based on truncated Schwinger-Dyson equations [133–138] while another approach uses renormalization group methods [139]. In many supersymmetric theories, the boundary of the conformal window can be determined, as discussed in [140]. There are also estimates for non-supersymmetric models based on ideas from supersymmetry [141]. Introductory reviews of lattice gauge theory research in this area can be found in [142–145]. Recent research includes the study of model with fermions in the fundamental representation [146–150], the adjoint representation [151–157], and the symmetric representation [158–162].

B. Conformality and duality in Abelian models

There is an interesting class of lattice models that share some of the features of gauge theories with conformal windows: $Z(N)$ spin systems in two dimensions and $Z(N)$ gauge

theories in four dimensions. Our understanding of these models originates in the two-dimensional XY model, which is a spin model with a global $U(1)$ symmetry. The order parameters of the model are the operators

$$S_p(x) = e^{ip\phi(x)} \quad (156)$$

where $\phi(x)$ takes on values between 0 and 2π . The XY model provides the principle example of a phase transition driven by topological excitations, the Berezinsky-Kosterlitz-Thouless (BKT) transition [163, 164]. Low-temperature arguments indicate that the XY model has a low-temperature gapless phase represented by a line of critical points, along which critical indices vary continuously with the temperature. On the other hand, high-temperature expansions indicate a gapped phase at high temperatures. The phase transition that separates the two phases is driven by vortices, configurations of spins that have a non-trivial winding number. In a continuum notation, these are configurations that have

$$\oint \nabla \phi \cdot dx = 2\pi n \quad (157)$$

where n is a non-zero integer. Vortices act as a two-dimensional classical Coulomb gas, interacting via a long-ranged logarithmic interaction. In the low-temperature phase, vortex pairs bind tightly to form bound states of zero vorticity and have no effect on the large-distance behavior. At the critical temperature, vortices unbind to form a classical Coulomb plasma, giving rise to a mass gap, the Debye mass of the plasma. From a naive continuum field theory point of view, the XY model has only massless spin-wave excitation, with an Euclidean action given by

$$S = \int d^2x \frac{J^2}{2} (\partial\phi)^2 \quad (158)$$

where J^2 is a coupling constant inversely proportional to the temperature. It is convenient to rescale the field ϕ by $\phi \rightarrow \phi/J$ so that the kinetic term of the action has conventional normalization. Vortices are created and destroyed by operators of the form

$$V_p(x) = e^{i2\pi p J \tilde{\phi}(x)} \quad (159)$$

where ϕ and $\tilde{\phi}$ are related by

$$i\partial_\mu \phi = \epsilon_{\mu\nu} \partial^\nu \tilde{\phi} \quad (160)$$

The effect of vortices gives rise to an effective action

$$S_{eff} = \int d^2x \left[\frac{J^2}{2} (\partial\phi)^2 - \sum_{p=1}^{\infty} 2y_p \Lambda^2 \cos(2\pi p J \tilde{\phi}) \right] \quad (161)$$




Figure 12: Relevance of V_1 for the sine-Gordon model.

where Λ is a cutoff on the order of the lattice spacing and y_p is a dimensionless activity for a vortex of winding number $\pm p$ [165, 166]. This generalized sine-Gordon model gives rise to Coulomb gas of vortices when the partition function is expanded in the activities. Generally speaking, it is only necessary to consider the $p = \pm 1$ vortices. This model can be perturbed to a model with $Z(p)$ symmetry by the addition of an operator $-h_p \Lambda^2 \cos(pJ^{-1}\phi)$ which explicitly breaks the $U(1)$ symmetry down to $Z(p)$:

$$S_{eff} = \int d^2x \left[\frac{J^2}{2} (\partial\phi)^2 - 2y_1 \Lambda^2 \cos(2\pi J\tilde{\phi}) - 2h_p \cos(p\phi/J) \right]. \quad (162)$$

This model has a duality under the interchange

$$\begin{aligned} \phi &\leftrightarrow \tilde{\phi} \\ h_p &\leftrightarrow y_1 \\ \frac{p}{J} &\leftrightarrow 2\pi J \end{aligned}$$

which is a generalization of Kramers-Wannier duality for the Ising model [167, 168].

A simple understanding of the renormalization-group behavior of the XY model is provided by analyzing the multiplicative renormalization of the spin-wave and vortex operators associated with normal ordering [169, 170]. For a massless free field, the relation between




Figure 13: Actual sine-Gordon flow.

the bare field $\exp(i\beta\phi)$ and the normal-ordered field $N[\exp(i\beta\phi)]$ is

$$\Lambda^2 e^{i\beta\phi} = \mu^2 \left(\frac{\Lambda}{\mu} \right)^{2 - \frac{\beta^2}{4\pi}} [e^{i\beta\phi}] \quad (163)$$

where μ is arbitrary. An interaction term of this type will be relevant if $\beta^2 < 8\pi$ and irrelevant if $\beta^2 > 8\pi$. When applied to the XY model, this implies that vortices are relevant only if $J^{-2} > \pi/2$, in the high-temperature phase. This leads to the simple picture shown in Fig. 12. Taking into account the renormalization of the vortex activity y_1 leads to a more complete picture of the renormalization group flow for the XY model, as shown in Fig. 13. Applying the same simple condition to the $Z(p)$ model indicates that the spin-wave symmetry breaking term $S_p + S_{-p}$ is relevant if $J^{-2} < 8\pi/p^2$, corresponding to low temperatures. If $p > 4$, there is a gap between the regions where neither spin-waves induced by $h_p \neq 0$ nor vortices are relevant, as shown in Fig. 14. In this intermediate conformal window, correlation functions decay algebraically; there is no mass gap. This phase structure is only part of a larger picture for more general $Z(p)$ lattice models. As we have seen in Sections III and IV, systems with $Z(p)$ symmetries can break spontaneously to a non-trivial subgroup of $Z(p)$. This also occurs in $d = 2$ $Z(p)$ spin systems [171–173]; for example $Z(6)$ can break spontaneously to $Z(2)$ or $Z(3)$. For $p = 2$ and $p = 3$, the regions of vortex and spin-wave relevance overlap, and must be handled as special cases.

These results extend to gauge theories in several different ways. The analysis of two dimensional systems by Jose *et al.* [165] can be extended to Abelian lattice gauge the-




Figure 14: Relevance of S_p and V_1 operators in a $Z(p)$ model for $p > 4$.

ories. In four-dimensional Abelian lattice models, the analogs of the point-like vortices of the two-dimensional spin models are closed magnetic monopole loops [174]. Similar to two-dimensional spin systems, four-dimensional $Z(p)$ lattice gauge theories have a duality between monopole loops and loops of charged particles. For $p > 4$, there is a gapless intermediate phase where neither the electric nor the magnetic operators are relevant [175]. In three-dimensional Abelian lattice models, the topological excitations are point-like, as is the case in Polyakov's treatment of the continuum three-dimensional $SU(2)$ gauge theory broken to $U(1)$. This is not accidental, because there is a clear relation between Abelian duality on the lattice and in the continuum. At finite temperature, *i.e.*, in a $R^2 \times S^1$ geometry, Polyakov's model can be analyzed in a manner analogous to $d = 2$ spin systems. Three-dimensional monopole solutions made periodic on S^1 play the role of vortices, and early work identified the phase transition to be in the XY-model, or $U(1)$, universality class [176]. As in 3+1 dimensions, adjoint Polyakov lines associated with $W^\pm$ gauge bosons ensure that the symmetry group of the low-energy theory remains $Z(2)$ [177].

Another interesting connection between the behavior of $d = 2$ spin models and $d = 4$ gauge theories is found in the suggestion by Kaplan et al. that there may be two nontrivial fixed points in gauge theories with a conformal window [178]. In this scenario, there is a UV fixed point at $g^2 = 0$, followed by a non-trivial infrared fixed point, followed by a non-trivial ultraviolet fixed point at a larger value of the g^2 . As N_f decreases, the two non-trivial fixed

points annihilate, which is to say they move off into the complex plane. This behavior would give rise to BKT scaling behavior in the region near where the fixed point merger occurs.

C. Relevance of topological objects on $R^3 \times S^1$

Unsal and Poppitz have proposed an alternative criterion for locating the boundary between confining and conformal gauge theories [179, 180]. Their work is based on the relevance or irrelevance of the topological objects relevant for confinement on $R^3 \times S^1$, analogous to the criterion for the relevance of vortices in $d = 2$ Abelian spin systems. For all center-symmetric gauge theories on $R^3 \times S^1$, the mass scale induced by topological excitations can be written in the form

$$m(L) \sim \frac{1}{L} \exp \left[-q \frac{8\pi^2}{g^2(L) N} \right] \quad (164)$$

where q is $\mathcal{O}(1)$ and depends on the details of the theory. With the use of the renormalization group, we can write this as

$$m(L) \sim \Lambda (\Lambda L)^{q\bar{b}_1-1} \quad (165)$$

where $\bar{b}_1 \equiv b_1/N$. Requiring that the theory remain in the center-symmetric, confined phase as L becomes large puts a constraint on the theory. In the limit $L \rightarrow \infty$, the topological excitations are relevant if $q\bar{b}_1 > 1$ and the theory is confined for large L . If $q\bar{b}_1 < 1$, the theory is not confined. While conceptually simple, the details of the calculations for different models are somewhat technical. The parameter q is an integer determined essentially by how many monopoles, each of action $8\pi^2/g^2 N$, are required in the process that generates the mass gap. For N_f^D flavors of Dirac fermions in the fundamental representation, the conformal window is estimated to lie in the range

$$\frac{5}{2}N \leq N_f^D \leq \frac{11}{2}N \quad (166)$$

while for N_f flavors of Weyl fermions in the adjoint representation the range is

$$4 \leq N_f^W \leq \frac{11}{2} \quad (167)$$

where $N_f^W = 2N_f^D$ if Dirac fermions are used. Estimates of the conformal window for other representations can also be made [179]. The estimates are comparable to estimates made using other methods, but the connection between this method and others is at present unclear.

VI. GAUGE THEORIES IN THE LARGE- N LIMIT

Many theories simplify in the large- N limit, where N is the dimension of some internal symmetry group such as $SU(N)$ taken to be large. Although physical QCD has $N = 3$, there is reason to believe the large- N limit provides a good description of many features [181, 182]. Lattice simulations at finite values of N have been able to provide accurate extrapolations to the large- N limit for some quantities [183, 184]. For example, lattice simulations of $SU(N)$ gauge theories in $(2 + 1)$ dimensions have provided evidence for $1/N$ corrections in string tension scaling laws [38]. The AdS/CFT correspondence, which relates gauge theories to gravity duals in the large- N limit [185], provides another reason for interest in large- N gauge theories; see [186, 187] for reviews.

The large- N limit began as an approximation in statistical mechanics [188], but was quickly applied to models with scalars ϕ^a [189] or fermions ψ^a [190] in the vector representation of the groups $O(N)$ or $SU(N)$. Such models are exactly solvable in the large- N limit. Field theories with particles in the adjoint representation, such as gauge bosons, are much more difficult.

The original approach of 't Hooft to the large- N limit in $SU(N)$ Yang-Mills theories was based around a detailed analysis of classes of Feynman diagrams using an ingenious double line notation [181]. This notation is based on the group properties of $SU(N)$ propagators: for fundamental representation fermions, we have

$$\langle \psi^a(x) \bar{\psi}_b(y) \rangle = \delta_b^a S(x - y) \quad (168)$$

but for gauge bosons, which are in the adjoint representation, we have

$$\langle A_{\mu b}^a(x) A_{\nu d}^c(y) \rangle = \left[\delta_d^a \delta_b^c - \frac{1}{N} \delta_b^a \delta_d^c \right] D_{\mu\nu}(x - y) \quad (169)$$

In this notation, quarks in N , the fundamental representation of $SU(N)$, and antiquarks in $\bar{N}$ are represented by single lines with arrows. Gauge bosons in the adjoint representation are represented by a double line with one arrow in each direction; other representations may be represented by other combinations of arrows.

The Euclidean Lagrangian for an $SU(N)$ gauge theory can be written as

$$L_E = \frac{1}{2g^2} \text{Tr}_F [F_{\mu\nu}^2] = \frac{N}{2\lambda} \text{Tr}_F [F_{\mu\nu}^2] \quad (170)$$

we have introduced $\lambda \equiv g^2 N$, which will be taken to be finite in a limit where $g \rightarrow 0$ as $N \rightarrow \infty$. It is obvious that every gauge boson propagator will carry a factor of $1/N$, and the cubic and quartic vertices will carry a factor of N . Fermion fields can be rescaled by $\psi \rightarrow N^{1/2}\psi$ so that similar properties hold.

Any vacuum diagram can be regarded as a polygon, where every closed loop can be regarded as a face. A vacuum diagram can then be characterized by its number of faces F , number of vertices V and number of edges E . Each face generates a factor of N from a trace. Every edge is a propagator, and carries with a factor of N^{-1} while every vertex carries a factor of N . Thus each vacuum graph carries a factor of

$$N^{F-E+V} \quad (171)$$

However, $F - E + V$ is the Euler number χ_E of the surface, a topological invariant. It may be calculated from the number of handles H of a surface and the number of holes B (for boundary) as

$$\chi_E = 2 - 2H - B. \quad (172)$$

Thus the maximum power of N associated with a vacuum diagram is N^2 , for diagrams with $H = B = 0$. Because each fundamental representation loop gives rise to a hole, fundamental representation particles begin contributing at order N . This behavior is what we see in the perturbative calculation of the effective potential for the Polyakov effective potential. The effects of quarks are suppressed by a factor N_f/N relative to gluons. It should be noted that there is another interesting large- N limit, the Veneziano limit, in which N_f is taken to infinity at the same time as N , with the ratio $x = N_f/N$ fixed [191]. This generates a large- N expansion where particles in the fundamental representation are not suppressed.

A key feature is large- N factorization. Suppose, for example, that O_1 and O_2 are operators transforming according to the adjoint representation of the group. Then $Tr O_i/N$ has a smooth limit as $N \rightarrow \infty$. Large- N factorization states that in the large- N limit

$$\left\langle \frac{1}{N} Tr O_1 \frac{1}{N} Tr O_2 \right\rangle \rightarrow \left\langle \frac{1}{N} Tr O_1 \right\rangle \left\langle \frac{1}{N} Tr O_2 \right\rangle + O(1/N^2) \quad (173)$$

up to terms of $O(1/N^2)$. For Wilson loops, large- N factorization leads to a closed, exact equation in the $N \rightarrow \infty$ limit, the Migdal-Makeenko equation [192]. It was quickly realized that many aspects are better captured by postulating the existence of a large- N saddle point of some large- N effective action analogous to the gap equation in vector-like models. This

saddle point, really a gauge orbit in the space of fields, is often referred to as the master field [193]. While it is possible to find the master field for some model systems in lower dimensions with fields in the adjoint representation [194], the problem of determining the master field for four-dimensional gauge theories has not been solved.

In recent years, much of the work on gauge theories in the large- N limit has employed gauge-string duality to explore strong-coupling behavior using generalizations of the AdS/CFT correspondence [185]. For a review emphasizing the relation of gauge/string duality to finite temperature QCD, see [187]. Attempts to model the phase structure of QCD based on the AdS/QCD approach [195] may be found in [196–199].

A. Eguchi-Kawai models

A direct attempt at constructing a large- N reduction for lattice gauge theories was first made in 1982 by Eguchi and Kawai [200]. They defined a lattice gauge theory on a single lattice site and showed that Wilson loops in an infinite volume theory could also be obtained from their reduced model in the large- N limit if two conditions were met: large- N factorization and center symmetry must both hold in all directions. This latter restriction implies that the reduction can only hold in the confined phase.

The original Eguchi-Kawai (EK) model may be derived by observing that the master field must be translation invariant. For a lattice gauge theory, this means that the value of the link variable at every site must be a gauge transform of the field at the origin

$$U_\mu(x) \rightarrow D^\dagger(x) \tilde{U}_\mu D(x + \mu) \quad (174)$$

where $\tilde{U}_\mu$ denotes the link variables at the origin. The matrix $D(x)$ can be taken to be a diagonal matrix of the form

$$D(x) = \exp(-ix \cdot P) \quad (175)$$

where P_μ is a diagonal Hermitian matrix of the form $P_\mu = \text{diag}(p_{1\mu}, \dots, p_{N\mu})$. Gauge-invariant quantities immediately collapse. For example, a rectangular $R \times T$ Wilson loop in the $\mu - \nu$ plane becomes

$$\left\langle \text{Tr} \left(\tilde{U}_\mu \right)^T \left(\tilde{U}_\nu \right)^R \left(\tilde{U}_\mu^\dagger \right)^T \left(\tilde{U}_\nu^\dagger \right)^R \right\rangle \quad (176)$$

and the Wilson action collapses to

$$S_{EK} = \mathcal{VT} \sum_{\mu > \nu} \frac{\beta}{N} \text{Re Tr} \left[\tilde{U}_\mu \tilde{U}_\nu \tilde{U}_\mu^\dagger \tilde{U}_\nu^\dagger \right] \quad (177)$$

where $\mathcal{VT}$ is the volume of space-time. Note that global center symmetry transformations of the form $\tilde{U}_\mu \rightarrow z_\mu \tilde{U}_\mu$ is a symmetry of S_{EK} .

In the infinite volume gauge theory, the expectation value of non-closed paths such as

$$\left\langle \text{Tr} \left(\tilde{U}_\mu \right)^T \right\rangle \quad (178)$$

is zero as a consequence of gauge-invariance and Elitzur's theorem. In the Eguchi-Kawai model, this expectation value will vanish if center symmetry is unbroken, but need not vanish if center symmetry is broken. A detailed comparison of the Schwinger-Dyson equations for both theories confirms that they are equivalent only if center symmetry is unbroken. The similarity with the case of finite temperature immediately suggests that center symmetry will be broken in the large- N limit, and the equivalence will fail. This turns out to be the case [201–203]. Several attempts have been made to modify the original EK model to maintain center symmetry. The first was the Quenched EK (QEK) model, in which the eigenvalues of the link variables are treated as quenched variables, forcing the link variables to maintain center symmetry [201]. While early lattice simulation did indicate that the QEK model was viable, but recent lattice simulations provides strong evidence for the breakdown of the reduction [204]. Another variant of the original model that originally seemed promising is the Twisted EK (TEK) model [205], but more recent extensive lattice simulations indicate center symmetry breaking in this model as well [206–208]. An interesting alternative to single-site reduction is to simulate large- N theories on lattices of size N_s^4 , and keep N_s sufficiently large that the deconfinement transition is avoided [184, 209]. As long as center symmetry is maintained, such models will be equivalent to infinite-volume theories in the limit of infinite N . This behavior, a hallmark of a successful large- N reduction, is often generally described as volume independence.

B. Planar equivalence

The most prominent example of planar equivalence is the equivalence, at large N , of the $\mathcal{N} = 1$ super-Yang-Mills model to an $SU(N)$ gauge theory with fermions in either the

symmetric or the antisymmetric two-index representation. In other words, in the large- N limit, $SU(N)$ gauge theories with fermions in the adjoint representation (Adj) are equivalent to the same gauge theory with fermions in either the symmetric (S) or antisymmetric (A) representation. The history of the subject has an interesting arc; see [210] for an introductory discussion. There is a very general formalism for describing many forms of large- N equivalences [211], with which a detailed discussion of many forms of planar equivalence can be carried out [212–214]. For our purposes, the equivalence of $SU(N)_{Adj,A,S}$ can be understood from the equivalence of the loop equations for the different theories [215]. Supersymmetric gauge theories have Majorana fermions in the adjoint representation which are superpartners of the gauge bosons. While the effects of fundamental representation fermions are suppressed in the large- N limit, the effects of adjoint representation fermions are not, because the number of components in the representation grows as N^2 in the large- N limit. In general, the dimensionality d_R of a given representation R is easily determined from its character $\chi_R(U) \equiv \text{Tr}_R U$ via $d_R = \chi_R(I)$. For example, we have for the adjoint representation

$$\chi_{Adj}(U) = \chi_F(U) \chi_{\bar{F}}(U) - 1 \quad (179)$$

so $d_{Adj} = N^2 - 1$. The characters of the symmetric and antisymmetric representations $\chi_{S/A}(U)$ are formed from the product of two characters in the fundamental representation and given by

$$\chi_{S/A}(U) = \frac{1}{2} [\chi_F^2(U) \pm \chi_F(U)] \quad (180)$$

with dimensions $N(N \pm 1)/2$. These representations are sometimes referred to as bifermion representations. Because the number of fields grows as N^2 , we expect the effects of internal loops of S/A fermions to survive in the large- N limit in the same way that adjoint fermion loops do.

Let us imagine integrating out the fermions in an $SU(N)$ gauge theory with either adjoint, symmetric or antisymmetric fermions [216]. In all cases, the logarithm of the fermion determinant can be expressed as a sum of Wilson loops in R^4 ; in a geometry like $R^3 \times S^1$ or T^4 , the sum includes Polyakov loops as well as Wilson loops. In the large- N limit, factorization tells us that on R^4 we have

$$2 \langle \chi_{S/A}(W) \rangle \sim \langle \chi_F(W) \rangle \langle \chi_F(W) \rangle \sim \langle \chi_F(W) \rangle \langle \chi_F^*(W) \rangle \sim \langle \chi_{Adj}(W) \rangle \quad (181)$$

where the expectation value is over the gauge fields. The apparent mis-match of a factor

of two disappears in the functional determinants because the fermions in the symmetric or antisymmetric representations are Dirac fermions, while the adjoint fermions are Majorana fermions with half the degrees of freedom. A proof of the equivalence at the level of Feynman diagrams can also be given using 't Hooft's double line notation [217]. In the confined phase on T^4 or a similar geometry, all the expectation value for the Polyakov loops are zero

$$2\langle\chi_{S/A}(P)\rangle\sim\langle\chi_F(P)\rangle\langle\chi_F(P)\rangle\sim\langle\chi_F(P)\rangle\langle\chi_F^*(P)\rangle\sim\langle\chi_{Adj}(P)\rangle=0\quad(182)$$

and any of the reduced gauge theories that is confining is completely equivalent to the theories on R^4 , realizing volume independence. However, this equivalence fails in non-confining phases, because the N -ality of the symmetric and antisymmetric representations is two, but for the adjoint representation it is zero. Thus we return once again to the issue of maintaining confinement in small volumes.

C. Confinement for finite volume

As we have seen in section IV, suitable modifications of the action can lead to a confining phase on $R^3 \times S^1$ for small circumference L . It is natural to look at these methods for the possible construction of useful reduced large- N models. However, the confining phase is only one of many possible phases as N becomes large [48, 49]. In principle, it is possible to modify the large- N gauge action on T^4 by adding double-trace deformations

$$S \rightarrow S + \sum_{k=1}^{[N/2]} \sum_{\mu} a_k \int d^4x \text{Tr}_A P_{\mu}(x)^k. \quad(183)$$

where $P_{\mu}(x)$ is the Polyakov loop in the direction μ based at the spacetime point x . There is a harmless redundancy induced by the integration over x , because all points a given loop path are integrated over. For a single-site lattice action, the corresponding modification is

$$S \rightarrow S + \sum_{k=1}^{[N/2]} \sum_{\mu} a_k \text{Tr}_A U_{\mu}^k. \quad(184)$$

because the single-site link variable is the Polyakov loop. If the coefficients a_k are sufficiently large, then the system will remain in the confined phase, with $\langle\text{Tr}_A P^k\rangle = 0$. In the confined phase, the precise values of the coefficients don't matter; we would have a self-consistent finite-volume reduction of the pure gauge theory [93].

Adjoint fermions with periodic boundary conditions offer another possible route to the construction of a large- N finite-volume reduction [97, 98, 119, 218–226]. This model is also of direct phenomenological interest. In physical QCD with $N = 3$, we normally think of quarks in the fundamental representation as playing no role in the large- N limit. However, antiquarks in physical QCD are in the $\bar{3}$ representation, which is an antisymmetric representation. In this sense, physical QCD has two different large- N limits. The large- N limit, an $SU(N)$ gauge theory with N_f Dirac fermions in the adjoint representation is equivalent to the theory with $2N_f$ Dirac fermions in the antisymmetric representation, which is QCD with $2N_f$ flavors when $N = 3$. Analytic calculations of the lattice form of the one-loop effective potential [97, 119, 218, 221, 222] indicate a rich phase structure as the fermion mass is varied, similar to that found in the continuum on $R^3 \times S^1$ [48, 119]. However, it should be kept in mind that chiral symmetry breaking effects, which affect Polyakov loop terms in the effective potential, are generally not included in these calculations; at present, these effects can only be estimated using phenomenological models [71]. Thus lattice simulations are essential not only to determine the properties of the confined phase, but also to confirm that it exists. The key issue is finding a confined phase in lattice simulations that survives the extrapolation to the continuum limit where the bare lattice coupling g^2 goes to zero. Of course, the success of the large- N reduction may depend on the lattice fermion implementation. In the single-site model, analytic calculations indicate that center symmetry is spontaneously broken with naive fermions but unbroken with overlap fermions [223]. Lattice simulations of $SU(3)$ with 2 flavors of staggered fermions in the adjoint representation on $R^3 \times S^1$ show that a confined phase does exist for sufficiently small fermion mass [96]. Simulations of the single-site theory using Wilson fermions also show a region in the parameter space of fermion mass and lattice coupling where center symmetry is unbroken and an extrapolation to the continuum limit appears plausible [98, 225]; see also [224]. Simulations with overlap fermions [226] also indicated a confined region. The most recent lattice results [227] of a single-site model with two flavors of Wilson fermions show a center-symmetric region for N as large as 53 and 't Hooft coupling $g^2 N$ as small as 0.005, so the prospects for a successful large- N reduction appear bright.

VII. CONCLUSIONS

We have entered a new phase in our understanding of the phase structure of gauge theories. The phase structure of gauge theories at finite temperature was known to be rich. By itself, however, temperature alone does not give us a large window where continuum analytical results and lattice simulations are both useful. We have now a class of theories that can be successfully studied using both analytical methods and lattice simulations, built on our understanding of finite temperature physics. On $R^3 \times S^1$, the use of double-trace deformations or periodic adjoint fermions allows us to study confinement in a region where semiclassical methods are valid, and check our results with lattice simulations. At the same time, we have access to many new phases, most of which are partially-confining phases. Nor are we restricted to an $R^3 \times S^1$ geometry; other geometries are available, and largely unexplored. By extending finite-temperature physics to this larger class of models, we have seen how instantons and non-Abelian monopoles can play a role in confinement, realizing some long-held ideas about the nature of the confining phase. At the same time, we have seen how topological effects can survive in the large- N limit. We also have a new, promising class of single-site large- N models to explore.

There is more to be done. Most of the lattice simulations to date have only checked analytic predictions of overall phase structure, and predictions for, *e.g.*, the behavior of string tensions have not been checked in simulations. Although semiclassical techniques have a natural range of validity on $R^3 \times S^1$ given by $N\Lambda L \ll 1$, we do not know the general validity of the concepts useful for small L . We know that a double-trace deformation connects the small- L confining phase to the large- L confining phase without an intervening phase transition, but we do not know how physical quantities change along a path that connects the two. We do not yet have a complete understanding of string-tension scaling laws, even in the small- L region, and there are many other questions that remain to be asked and answered.

Acknowledgments

The author would like to thank his collaborators Peter Meisinger, Joyce Myers, and Hiro Nishimura for sharing their insights into this subject, Mithat Unsal for many stimulating

discussions, and his colleagues Mark Alford and Francesc Ferrer for their helpful advice. The author gratefully acknowledges the support of this work by the U.S. Dept. of Energy under Grant 91ER40628.

-
- [1] M. Unsal, Phys.Rev.Lett. **100**, 032005 (2008), 0708.1772.
 - [2] J. C. Myers and M. C. Ogilvie, Phys.Rev. **D77**, 125030 (2008), 0707.1869.
 - [3] P. M. Chaikin and T. C. Lubensky, *Principles of Condensed Matter Physics* (Cambridge University Press, Cambridge, 1995).
 - [4] L. P. Kadanoff, W. Gotze, D. Hamblen, R. Hecht, E. Lewis, et al., Rev.Mod.Phys. **39**, 395 (1967).
 - [5] K. Wilson and J. B. Kogut, Phys.Rept. **12**, 75 (1974).
 - [6] K. G. Wilson, Phys.Rev. **B4**, 3174 (1971).
 - [7] K. G. Wilson, Phys.Rev. **B4**, 3184 (1971).
 - [8] K. G. Wilson and M. E. Fisher, Phys.Rev.Lett. **28**, 240 (1972).
 - [9] L. Dolan and R. Jackiw, Phys.Rev. **D9**, 3320 (1974).
 - [10] S. Weinberg, Phys.Rev. **D9**, 3357 (1974).
 - [11] W. Selke, Phys. Rep. **170**, 213 (1988).
 - [12] S. Ostlund, Phys.Rev. **B24**, 398 (1981).
 - [13] S. Howes, L. Kadanoff, and M. Den Nijs, Nucl.Phys. **B215**, 169 (1983).
 - [14] M. Ogilvie and P. Meisinger, SIGMA **5**, 047, 14 p. (2009).
 - [15] M. Ogilvie, P. Meisinger, and T. Wiser, International Journal of Theoretical Physics **50**, 1042 (2011).
 - [16] M. G. Alford, J. A. Bowers, and K. Rajagopal, Phys.Rev. **D63**, 074016 (2001), hep-ph/0008208.
 - [17] A. Larkin and Y. Ovchinnikov, Zh.Eksp.Teor.Fiz. **47**, 1136 (1964).
 - [18] P. Fulde and R. A. Ferrell, Phys.Rev. **135**, A550 (1964).
 - [19] J. Langer, Annals Phys. **41**, 108 (1967).
 - [20] S. R. Coleman, Phys.Rev. **D15**, 2929 (1977).
 - [21] C. G. Callan and S. R. Coleman, Phys.Rev. **D16**, 1762 (1977).
 - [22] E. J. Weinberg and A.-q. Wu, Phys.Rev. **D36**, 2474 (1987).

- [23] E. W. Kolb and M. S. Turner, *The Early Universe*, Frontiers in Physics (Westview Press, Boulder, CO, 1990).
- [24] E. Kolb, Edward W. and E. Turner, Michael S., *The Early Universe: Reprints* (Addison-Wesley Pub. Co., Advanced Book Program, 1988).
- [25] K. G. Wilson, Phys.Rev. **D10**, 2445 (1974).
- [26] M. Luscher, Nucl.Phys. **B180**, 317 (1981).
- [27] M. Luscher and P. Weisz, JHEP **0207**, 049 (2002), hep-lat/0207003.
- [28] H. B. Meyer, JHEP **0605**, 066 (2006), hep-th/0602281.
- [29] O. Aharony and E. Karzbrun, JHEP **0906**, 012 (2009), 0903.1927.
- [30] O. Aharony and M. Field, JHEP **1101**, 065 (2011), 1008.2636.
- [31] F. Gliozzi, M. Pepe, and U.-J. Wiese, Phys.Rev.Lett. **104**, 232001 (2010), 1002.4888.
- [32] F. Gliozzi, M. Pepe, and U.-J. Wiese, JHEP **1101**, 057 (2011), 1010.1373.
- [33] M. Pepe, PoS **LATTICE2010**, 017 (2010), 1011.0056.
- [34] J. Greensite, Prog.Part.Nucl.Phys. **51**, 1 (2003), hep-lat/0301023.
- [35] M. R. Douglas and S. H. Shenker, Nucl.Phys. **B447**, 271 (1995), hep-th/9503163.
- [36] A. Hanany, M. J. Strassler, and A. Zaffaroni, Nucl.Phys. **B513**, 87 (1998), hep-th/9707244.
- [37] A. Armoni and M. Shifman, Nucl.Phys. **B671**, 67 (2003), hep-th/0307020.
- [38] B. Bringoltz and M. Teper, Phys.Lett. **B663**, 429 (2008), 0802.1490.
- [39] J. Greensite, B. Lucini, and A. Patella, Phys.Rev. **D83**, 125019 (2011), 1101.5344.
- [40] S. Deldar, Phys.Rev. **D62**, 034509 (2000), hep-lat/9911008.
- [41] G. S. Bali, Phys.Rev. **D62**, 114503 (2000), hep-lat/0006022.
- [42] P. N. Meisinger, T. R. Miller, and M. C. Ogilvie, Phys.Rev. **D65**, 034009 (2002), hep-ph/0108009.
- [43] M. Billo, M. Caselle, A. D’Adda, and S. Panzeri, Int.J.Mod.Phys. **A12**, 1783 (1997), hep-th/9610144.
- [44] M. Blume, V. J. Emery, and R. B. Griffiths, Phys. Rev. A **4**, 1071 (1971).
- [45] I. Halliday and A. Schwimmer, Phys.Lett. **B101**, 327 (1981).
- [46] I. Halliday and A. Schwimmer, Phys.Lett. **B102**, 337 (1981).
- [47] J.-M. Drouffe and J.-B. Zuber, Phys.Rept. **102**, 1 (1983).
- [48] J. C. Myers and M. C. Ogilvie, JHEP **0907**, 095 (2009), 0903.4638.
- [49] P. N. Meisinger and M. C. Ogilvie, Phys. Rev. **D81**, 025012 (2010), 0905.3577.

- [50] K. Holland, P. Minkowski, M. Pepe, and U. Wiese, Nucl.Phys. **B668**, 207 (2003), hep-lat/0302023.
- [51] M. Pepe, PoS **LAT2005**, 017 (2006), hep-lat/0510013.
- [52] D. J. Gross, R. D. Pisarski, and L. G. Yaffe, Rev. Mod. Phys. **53**, 43 (1981).
- [53] N. Weiss, Phys. Rev. **D24**, 475 (1981).
- [54] J. Polonyi and K. Szlachanyi, Phys.Lett. **B110**, 395 (1982).
- [55] M. Ogilvie, Phys.Rev.Lett. **52**, 1369 (1984).
- [56] F. Green and F. Karsch, Nucl.Phys. **B238**, 297 (1984).
- [57] M. Gross and J. Wheeler, Nucl.Phys. **B240**, 253 (1984).
- [58] J. Langelage, S. Lottini, and O. Philipsen, JHEP **1102**, 057 (2011), 1010.0951.
- [59] B. Svetitsky and L. G. Yaffe, Nucl. Phys. **B210**, 423 (1982).
- [60] B. Lucini, M. Teper, and U. Wenger, Phys.Lett. **B545**, 197 (2002), hep-lat/0206029.
- [61] B. Lucini, M. Teper, and U. Wenger, JHEP **0401**, 061 (2004), hep-lat/0307017.
- [62] F. Karsch, Lect.Notes Phys. **583**, 209 (2002), hep-lat/0106019.
- [63] A. D. Linde, Phys.Lett. **B96**, 289 (1980).
- [64] K. Kajantie, M. Laine, K. Rummukainen, and Y. Schroder, Phys.Rev. **D67**, 105008 (2003), hep-ph/0211321.
- [65] G. Boyd, J. Engels, F. Karsch, E. Laermann, C. Legeland, et al., Nucl.Phys. **B469**, 419 (1996), hep-lat/9602007.
- [66] A. Papa, Nucl.Phys. **B478**, 335 (1996), hep-lat/9605004.
- [67] B. Beinlich, F. Karsch, E. Laermann, and A. Peikert, Eur.Phys.J. **C6**, 133 (1999), hep-lat/9707023.
- [68] M. Okamoto et al. (CP-PACS Collaboration), Phys.Rev. **D60**, 094510 (1999), hep-lat/9905005.
- [69] A. Dumitru, Y. Guo, Y. Hidaka, C. P. K. Altes, and R. D. Pisarski, Phys.Rev. **D83**, 034022 (2011), 1011.3820.
- [70] A. Dumitru, Y. Guo, Y. Hidaka, C. P. K. Altes, and R. D. Pisarski (2012), 1205.0137.
- [71] H. Nishimura and M. C. Ogilvie, Phys.Rev. **D81**, 014018 (2010), 0911.2696.
- [72] S. Klevansky, Rev.Mod.Phys. **64**, 649 (1992).
- [73] T. Hatsuda and T. Kunihiro, Phys.Rept. **247**, 221 (1994), hep-ph/9401310.
- [74] P. N. Meisinger and M. C. Ogilvie, Phys. Rev. **D65**, 056013 (2002), hep-ph/0108026.

- [75] J. I. Kapusta and C. Gale, *Finite-temperature field theory. Principle and applications. 2nd ed.* (Cambridge Monographs on Mathematical Physics. Cambridge: Cambridge University Press., 2006).
- [76] A. Hasenfratz and T. A. DeGrand, Phys.Rev. **D49**, 466 (1994), hep-lat/9304001.
- [77] P. N. Meisinger and M. C. Ogilvie, Phys.Rev. **D52**, 3024 (1995), hep-lat/9502003.
- [78] K. Kashiwa, R. D. Pisarski, and V. V. Skokov (2012), 1205.0545.
- [79] A. Gocksch and M. Ogilvie, Phys.Rev. **D31**, 877 (1985).
- [80] K. Fukushima, Phys.Lett. **B591**, 277 (2004), hep-ph/0310121.
- [81] A. Mocsy, F. Sannino, and K. Tuominen, Phys.Rev.Lett. **92**, 182302 (2004), hep-ph/0308135.
- [82] F. Karsch and M. Lutgemeier, Nucl.Phys. **B550**, 449 (1999), hep-lat/9812023.
- [83] J. Engels, S. Holtmann, and T. Schulze, Nucl.Phys. **B724**, 357 (2005), hep-lat/0505008.
- [84] P. de Forcrand and O. Jahn (0300), hep-lat/0503041.
- [85] O. Philipsen, PoS **LAT2005**, 016 (2006), hep-lat/0510077.
- [86] E. Loh, J. Gubernatis, R. Scalettar, S. White, D. Scalapino, et al., Phys.Rev. **B41**, 9301 (1990).
- [87] K. Fukushima and T. Hatsuda, Rept.Prog.Phys. **74**, 014001 (2011), 1005.4814.
- [88] S. Mandelstam, Phys.Lett. **B53**, 476 (1975).
- [89] S. Mandelstam, Phys.Rept. **23**, 245 (1976).
- [90] G. 't Hooft, Nucl.Phys. **B138**, 1 (1978).
- [91] G. 't Hooft, Nucl.Phys. **B153**, 141 (1979).
- [92] M. C. Ogilvie, P. N. Meisinger, and J. C. Myers (2007), arXiv:0710.0649 [hep-lat].
- [93] M. Unsal and L. G. Yaffe, Phys.Rev. **D78**, 065035 (2008), 0803.0344.
- [94] N. M. Davies, T. J. Hollowood, V. V. Khoze, and M. P. Mattis, Nucl. Phys. **B559**, 123 (1999), hep-th/9905015.
- [95] N. M. Davies, T. J. Hollowood, and V. V. Khoze, J. Math. Phys. **44**, 3640 (2003), hep-th/0006011.
- [96] G. Cossu and M. D'Elia, JHEP **0907**, 048 (2009), 0904.1353.
- [97] B. Bringoltz, JHEP **0906**, 091 (2009), 0905.2406.
- [98] B. Bringoltz and S. R. Sharpe, Phys.Rev. **D80**, 065031 (2009), 0906.3538.
- [99] A. M. Polyakov, Nucl.Phys. **B120**, 429 (1977).
- [100] K.-M. Lee and C.-h. Lu, Phys.Rev. **D58**, 025011 (1998), hep-th/9802108.

- [101] T. C. Kraan and P. van Baal, Phys.Lett. **B428**, 268 (1998), hep-th/9802049.
- [102] T. C. Kraan and P. van Baal, Nucl.Phys. **B533**, 627 (1998), hep-th/9805168.
- [103] M. Prasad and C. M. Sommerfield, Phys.Rev.Lett. **35**, 760 (1975).
- [104] E. Bogomolny, Sov.J.Nucl.Phys. **24**, 449 (1976).
- [105] T. W. Kirkman and C. K. Zachos, Phys.Rev. **D24**, 999 (1981).
- [106] E. Witten, Nucl.Phys. **B149**, 285 (1979).
- [107] C. E. Detar, Phys. Rev. **D32**, 276 (1985).
- [108] T. A. DeGrand and C. E. DeTar, Phys. Rev. **D34**, 2469 (1986).
- [109] G. S. Bali, J. Fingberg, U. M. Heller, F. Karsch, and K. Schilling, Phys. Rev. Lett. **71**, 3059 (1993), hep-lat/9306024.
- [110] F. Karsch, E. Laermann, and M. Lutgemeier, Phys. Lett. **B346**, 94 (1995), hep-lat/9411020.
- [111] K. Zarembo, Nucl.Phys. **B463**, 73 (1996), hep-th/9510031.
- [112] M. Unsal, Phys.Rev. **D80**, 065001 (2009), 0709.3269.
- [113] E. Poppitz, T. Schaefer, and M. Unsal (2012), 1205.0290.
- [114] P. N. Meisinger and M. C. Ogilvie, Nucl.Phys.Proc.Suppl. **140**, 650 (2005), hep-lat/0409136.
- [115] O. Philipsen and H. Wittig, Phys.Rev.Lett. **81**, 4056 (1998), hep-lat/9807020.
- [116] F. Knechtli and R. Sommer (ALPHA collaboration), Phys.Lett. **B440**, 345 (1998), hep-lat/9807022.
- [117] F. Knechtli and R. Sommer (ALPHA Collaboration), Nucl.Phys. **B590**, 309 (2000), hep-lat/0005021.
- [118] M. Pepe and U.-J. Wiese, Phys.Rev.Lett. **102**, 191601 (2009), 0901.2510.
- [119] T. J. Hollowood and J. C. Myers, JHEP **0911**, 008 (2009), 0907.3665.
- [120] S. Hands, T. J. Hollowood, and J. C. Myers, JHEP **1007**, 086 (2010), 1003.5813.
- [121] S. Hands, T. J. Hollowood, and J. C. Myers, JHEP **1012**, 057 (2010), 1010.0790.
- [122] T. J. Hollowood, S. P. Kumar, and J. C. Myers, JHEP **1111**, 138 (2011), 1110.0696.
- [123] D. Simic and M. Unsal, Phys.Rev. **D85**, 105027 (2012), 1010.5515.
- [124] M. M. Anber, E. Poppitz, and M. Unsal, JHEP **1204**, 040 (2012), 1112.6389.
- [125] H. Nishimura and M. C. Ogilvie, AIP Conf. Proc. **1343**, 167 (2011), 1012.0333.
- [126] H. Nishimura and M. C. Ogilvie, Phys.Rev. **D85**, 065021 (2012), 1111.6101.
- [127] S. Elitzur, Phys.Rev. **D12**, 3978 (1975).
- [128] E. H. Fradkin and S. H. Shenker, Phys.Rev. **D19**, 3682 (1979).

- [129] N. Seiberg and E. Witten, Nucl. Phys. **B426**, 19 (1994), hep-th/9407087.
- [130] E. Poppitz and M. Unsal, JHEP **07**, 082 (2011), 1105.3969.
- [131] T. Banks and A. Zaks, Nucl.Phys. **B196**, 189 (1982).
- [132] F. Sannino, Acta Phys.Polon. **B40**, 3533 (2009), 0911.0931.
- [133] T. Appelquist, K. D. Lane, and U. Mahanta, Phys.Rev.Lett. **61**, 1553 (1988).
- [134] A. G. Cohen and H. Georgi, Nucl.Phys. **B314**, 7 (1989).
- [135] V. Miransky and K. Yamawaki, Phys.Rev. **D55**, 5051 (1997), hep-th/9611142.
- [136] T. Appelquist, J. Terning, and L. Wijewardhana, Phys.Rev.Lett. **77**, 1214 (1996), hep-ph/9602385.
- [137] T. Appelquist, A. Ratnaweera, J. Terning, and L. Wijewardhana, Phys.Rev. **D58**, 105017 (1998), hep-ph/9806472.
- [138] D. D. Dietrich and F. Sannino, Phys.Rev. **D75**, 085018 (2007), hep-ph/0611341.
- [139] H. Gies and J. Jaeckel, Eur.Phys.J. **C46**, 433 (2006), hep-ph/0507171.
- [140] K. A. Intriligator and N. Seiberg, Nucl.Phys.Proc.Suppl. **45BC**, 1 (1996), hep-th/9509066.
- [141] T. A. Ryttov and F. Sannino, Phys.Rev. **D76**, 105004 (2007), 0707.3166.
- [142] G. T. Fleming, PoS **LATTICE2008**, 021 (2008), 0812.2035.
- [143] T. DeGrand and A. Hasenfratz, Phys.Rev. **D80**, 034506 (2009), 0906.1976.
- [144] T. DeGrand (2010), 1010.4741.
- [145] L. Del Debbio (2011), 1102.4066.
- [146] A. Hasenfratz, Phys.Rev. **D80**, 034505 (2009), 0907.0919.
- [147] Z. Fodor, K. Holland, J. Kuti, D. Nogradi, and C. Schroeder, Phys.Lett. **B681**, 353 (2009), 0907.4562.
- [148] A. Hasenfratz, Phys.Rev. **D82**, 014506 (2010), 1004.1004.
- [149] Z. Fodor, K. Holland, J. Kuti, D. Nogradi, C. Schroeder, et al., Phys.Lett. **B703**, 348 (2011), 1104.3124.
- [150] T. Appelquist, G. Fleming, M. Lin, E. Neil, and D. Schaich, Phys.Rev. **D84**, 054501 (2011), 1106.2148.
- [151] L. Del Debbio, A. Patella, and C. Pica, Phys.Rev. **D81**, 094503 (2010), 0805.2058.
- [152] A. J. Hietanen, K. Rummukainen, and K. Tuominen, Phys.Rev. **D80**, 094504 (2009), 0904.0864.
- [153] L. Del Debbio, B. Lucini, A. Patella, C. Pica, and A. Rago, Phys.Rev. **D80**, 074507 (2009),

- 0907.3896.
- [154] F. Bursa, L. Del Debbio, L. Keegan, C. Pica, and T. Pickup, Phys.Rev. **D81**, 014505 (2010), 0910.4535.
 - [155] L. Del Debbio, B. Lucini, A. Patella, C. Pica, and A. Rago, Phys.Rev. **D82**, 014510 (2010), 1004.3206.
 - [156] H. S. Fukano and F. Sannino, Phys.Rev. **D82**, 035021 (2010), 1005.3340.
 - [157] T. DeGrand, Y. Shamir, and B. Svetitsky, Phys.Rev. **D83**, 074507 (2011), 1102.2843.
 - [158] Z. Fodor, K. Holland, J. Kuti, D. Nogradi, and C. Schroeder, JHEP **0911**, 103 (2009), 0908.2466.
 - [159] T. DeGrand, Phys.Rev. **D80**, 114507 (2009), 0910.3072.
 - [160] J. Kogut and D. Sinclair, Phys.Rev. **D81**, 114507 (2010), 1002.2988.
 - [161] T. DeGrand, Y. Shamir, and B. Svetitsky, Phys.Rev. **D82**, 054503 (2010), 1006.0707.
 - [162] J. Kogut and D. Sinclair, Phys.Rev. **D84**, 074504 (2011), 1105.3749.
 - [163] V. Berezinsky, Sov.Phys.JETP **32**, 493 (1971).
 - [164] J. Kosterlitz and D. Thouless, J.Phys.C **C6**, 1181 (1973).
 - [165] J. V. Jose, L. P. Kadanoff, S. Kirkpatrick, and D. R. Nelson, Phys.Rev. **B16**, 1217 (1977).
 - [166] M. C. Ogilvie, Annals Phys. **136**, 273 (1981).
 - [167] H. Kramers and G. Wannier, Phys.Rev. **60**, 252 (1941).
 - [168] H. Kramers and G. Wannier, Phys.Rev. **60**, 263 (1941).
 - [169] S. R. Coleman, Phys.Rev. **D11**, 2088 (1975).
 - [170] D. J. Amit, Y. Y. Goldschmidt, and G. Grinstein, J.Phys.A **A13**, 585 (1980).
 - [171] F. Alcaraz and R. Koberle, J.Phys.A **A13**, L153 (1980).
 - [172] F. Alcaraz and R. Koberle, J.Phys.A **A14**, 1169 (1981).
 - [173] P. Dorey, P. Provero, R. Tateo, and S. Vinti, J.Phys.A **A32**, L151 (1999), hep-th/9810202.
 - [174] T. Banks, R. Myerson, and J. B. Kogut, Nucl.Phys. **B129**, 493 (1977).
 - [175] A. Ukawa, P. Windey, and A. H. Guth, Phys.Rev. **D21**, 1013 (1980).
 - [176] N. O. Agasian and K. Zarembo, Phys.Rev. **D57**, 2475 (1998), hep-th/9708030.
 - [177] G. V. Dunne, I. I. Kogan, A. Kovner, and B. Tekin, JHEP **0101**, 032 (2001), hep-th/0010201.
 - [178] D. B. Kaplan, J.-W. Lee, D. T. Son, and M. A. Stephanov, Phys.Rev. **D80**, 125005 (2009), 0905.4752.
 - [179] E. Poppitz and M. Unsal, JHEP **0909**, 050 (2009), 0906.5156.

- [180] E. Poppitz and M. Unsal, JHEP **0912**, 011 (2009), 0910.1245.
- [181] G. 't Hooft, Nucl.Phys. **B72**, 461 (1974).
- [182] E. Witten, Nucl.Phys. **B160**, 57 (1979).
- [183] M. Teper, PoS **LAT2005**, 256 (2006), hep-lat/0509019.
- [184] R. Narayanan and H. Neuberger, PoS **LAT2007**, 020 (2007), 0710.0098.
- [185] J. M. Maldacena, Adv.Theor.Math.Phys. **2**, 231 (1998), hep-th/9711200.
- [186] K. Peeters and M. Zamaklar, Eur.Phys.J.ST **152**, 113 (2007), 0708.1502.
- [187] J. Casalderrey-Solana, H. Liu, D. Mateos, K. Rajagopal, and U. A. Wiedemann (2011), 1101.0618.
- [188] H. Stanley, Phys.Rev. **176**, 718 (1968).
- [189] S. R. Coleman, R. Jackiw, and H. D. Politzer, Phys.Rev. **D10**, 2491 (1974).
- [190] D. J. Gross and A. Neveu, Phys.Rev. **D10**, 3235 (1974).
- [191] G. Veneziano, Nucl.Phys. **B117**, 519 (1976).
- [192] Y. Makeenko and A. A. Migdal, Phys.Lett. **B88**, 135 (1979).
- [193] S. Coleman, *Aspects of Symmetry: Selected Erice Lectures* (Cambridge University Press, 1988).
- [194] E. Brezin, C. Itzykson, G. Parisi, and J. Zuber, Commun.Math.Phys. **59**, 35 (1978).
- [195] J. Erlich, E. Katz, D. T. Son, and M. A. Stephanov, Phys.Rev.Lett. **95**, 261602 (2005), hep-ph/0501128.
- [196] K. Kajantie, T. Tahkokallio, and J.-T. Yee, JHEP **0701**, 019 (2007), hep-ph/0609254.
- [197] C. Ballon Bayona, H. Boschi-Filho, N. R. Braga, and L. A. Pando Zayas, Phys.Rev. **D77**, 046002 (2008), 0705.1529.
- [198] U. Gursoy, E. Kiritsis, L. Mazzanti, and F. Nitti, Phys.Rev.Lett. **101**, 181601 (2008), 0804.0899.
- [199] U. Gursoy, E. Kiritsis, L. Mazzanti, and F. Nitti, Nucl.Phys. **B820**, 148 (2009), 0903.2859.
- [200] T. Eguchi and H. Kawai, Phys.Rev.Lett. **48**, 1063 (1982).
- [201] G. Bhanot, U. M. Heller, and H. Neuberger, Phys.Lett. **B113**, 47 (1982).
- [202] V. Kazakov and A. A. Migdal, Phys.Lett. **B116**, 423 (1982).
- [203] M. Okawa, Phys.Rev.Lett. **49**, 353 (1982).
- [204] B. Bringoltz and S. R. Sharpe, Phys.Rev. **D78**, 034507 (2008), 0805.2146.
- [205] A. Gonzalez-Arroyo and M. Okawa, Phys.Lett. **B120**, 174 (1983).

- [206] M. Teper and H. Vairinhos, Phys.Lett. **B652**, 359 (2007), hep-th/0612097.
- [207] T. Azeyanagi, M. Hanada, T. Hirata, and T. Ishikawa, JHEP **0801**, 025 (2008), 0711.1925.
- [208] W. Bietenholz, A. Bigarini, J. Nishimura, Y. Susaki, A. Torrielli, et al., PoS **LAT2007**, 049 (2007), 0708.1857.
- [209] J. Kiskis, R. Narayanan, and H. Neuberger, Phys.Rev. **D66**, 025019 (2002), hep-lat/0203005.
- [210] A. Armoni and M. Shifman (2007), hep-th/0702045.
- [211] P. Kovtun, M. Unsal, and L. G. Yaffe, JHEP **0507**, 008 (2005), hep-th/0411177.
- [212] P. Kovtun, M. Unsal, and L. G. Yaffe, Phys.Rev. **D72**, 105006 (2005), hep-th/0505075.
- [213] M. Unsal and L. G. Yaffe, Phys.Rev. **D74**, 105019 (2006), hep-th/0608180.
- [214] P. Kovtun, M. Unsal, and L. G. Yaffe, JHEP **0706**, 019 (2007), hep-th/0702021.
- [215] P. Kovtun, M. Unsal, and L. G. Yaffe, JHEP **0312**, 034 (2003), hep-th/0311098.
- [216] A. Armoni, M. Shifman, and G. Veneziano, Phys.Rev. **D71**, 045015 (2005), hep-th/0412203.
- [217] A. Armoni, M. Shifman, and G. Veneziano, Nucl.Phys. **B667**, 170 (2003), hep-th/0302163.
- [218] P. F. Bedaque, M. I. Buchoff, A. Cherman, and R. P. Springer, JHEP **0910**, 070 (2009), 0904.0277.
- [219] M. Hanada, L. Mannelli, and Y. Matsuo, Phys.Rev. **D80**, 125001 (2009), 0905.2995.
- [220] M. Hanada, L. Mannelli, and Y. Matsuo, JHEP **0911**, 087 (2009), 0907.4937.
- [221] B. Bringoltz, JHEP **1001**, 069 (2010), 0911.0352.
- [222] E. Poppitz and M. Unsal, JHEP **1001**, 098 (2010), 0911.0358.
- [223] A. Hietanen and R. Narayanan, JHEP **1001**, 079 (2010), 0911.2449.
- [224] T. Azeyanagi, M. Hanada, M. Unsal, and R. Yacoby, Phys.Rev. **D82**, 125013 (2010), 1006.0717.
- [225] S. Catterall, R. Galvez, and M. Unsal, JHEP **1008**, 010 (2010), 1006.2469.
- [226] A. Hietanen and R. Narayanan, Phys.Lett. **B698**, 171 (2011), 1011.2150.
- [227] B. Bringoltz, M. Koren, and S. R. Sharpe, Phys.Rev. **D85**, 094504 (2012), 1106.5538.